



Contribution ID: 159

Type: Talk

Roberge-Weiss transition in $N_f=2$ QCD with staggered fermions and $N_t=6$

Monday, July 25, 2016 5:05 PM (20 minutes)

The QCD phase diagram at imaginary chemical potential exhibits a rich structure and studying it can constrain the phase diagram at real values of the chemical potential. Moreover, at imaginary chemical potential standard numeric techniques based on importance sampling can be applied, since no sign problem is present.

In the last decade, a first understanding of the QCD phase diagram at imaginary chemical potential has been developed, but it is so far based on investigations on coarse lattices ($N_t = 4$, $a = 0.3$ fm). Considering the $N_f = 2$ case, at the Roberge-Weiss critical value of the imaginary chemical potential, the chiral/deconfinement transition is first order for light/heavy quark masses and second order for intermediate values of the mass: there are then two tricritical masses, whose position strongly depends on the lattice spacing and on the discretisation. On $N_t = 4$, we have the chiral $m_\pi^{\text{tric.}} \simeq 400$ MeV with unimproved staggered fermions and $m_\pi^{\text{tric.}} \simeq 900$ MeV with unimproved pure Wilson fermions.

Employing finite size scaling we investigate the change of this tricritical point between $N_t = 4$ and $N_t = 6$ as well as between Wilson and staggered discretisations.

Author: Mr SCIARRA, Alessandro (Goethe Universität)

Co-author: Prof. PHILIPSEN, Owe (Goethe Universität)

Presenter: Mr SCIARRA, Alessandro (Goethe Universität)

Session Classification: Nonzero Temperature and Density

Track Classification: Nonzero Temperature and Density