



Contribution ID: 81

Type: Talk

B_c decays from highly improved staggered quarks and NRQCD

Wednesday, July 27, 2016 9:40 AM (20 minutes)

We use both non-relativistic QCD (NRQCD) and fully relativistic formalisms to calculate semileptonic form factors for the decays $B_c \rightarrow \eta_c \ell \nu$ and $B_c \rightarrow J/\psi \ell \nu$ over the entire q^2 range. To achieve this we employ a highly improved lattice quark action at several lattice spacings down to $a=0.044$ fm, which allows a fully relativistic treatment of charm and simulation of the full q^2 range with controlled continuum extrapolation. We have two ways of treating the b quark: either with an $O(\alpha_s)$ improved NRQCD formalism or by extrapolating a heavy mass m_h to m_b in the relativistic formalism. Comparison of the two approaches provides an important cross-check of methodologies in lattice QCD. Nonperturbative renormalisation of the currents in the relativistic theory also allows us then to fix NRQCD-charm normalisation for b to c decays such as $B \rightarrow D$ and $B \rightarrow D^*$.

Primary author: Dr LYTLE, Andrew (University of Glasgow)

Co-authors: Dr COLQUHOUN, Brian (University of Glasgow); Prof. DAVIES, Christine (University of Glasgow); Dr MCNEILE, Craig (Plymouth University); Dr KOPONEN, Jonna (University of Glasgow)

Presenter: Dr LYTLE, Andrew (University of Glasgow)

Session Classification: Weak Decays and Matrix Elements

Track Classification: Weak Decays and Matrix Elements