

The quantum structure of EFT

Giampiero Passarino

Dipartimento di Fisica Teorica, Università di Torino, Italy
INFN, Sezione di Torino, Italy

The Final HiggsTools Meeting, Durham




constructing SMEFT

- Experiments occur at finite energy and measure $\mathbf{S}^{\text{eff}}(\Lambda)$
- Whatever QFT should give low energy $\mathbf{S}^{\text{eff}}(\Lambda)$, $\forall \Lambda < \infty$
- There is no fundamental scale above which $\mathbf{S}^{\text{eff}}(\Lambda)$ is not defined (K. Costello, Renormalization and EFT, AMS)
- $\mathbf{S}^{\text{eff}}(\Lambda)$ loses its predictive power if a process at $E = \Lambda$ requires ∞ renormalized parameters (J. Preskill, CALT-68-1493)

No phenomenology here, not even a discussion on the *validity* range of the SMEFT, i.e.

- SMEFT (linear representation) vs. HEFT (non-linear representation),
 - linear vs. quadratic implementation of $\dim = 6$ operators, i.e. positivity,
 - unitarity bounds
 - analyticity, etc.
- Waiting for the next SM or the next EFT, is SMEFT a consistent QFT from the point of view of removing UV poles (beware, an EFT is not strictly renormalizable)?

Note a/Λ^2 degeneracy



a fixed
 Λ small
is large MHOU

EFT is just the effective theory only designed for perturbative calculations of the S-matrix elements. Green functions may contain divergent contributions: we are only interested in Dyson-perturbation- scheme of calculating the S matrix.

Though the effective theory is renormalizable by construction it presents no interest until the problem of redundant operators is solved.

Thus it is necessary to point out an infinite number of renormalization prescriptions that allow one to fix the finite parts of counterterms. If this is done arbitrarily, the theory loses its predictive power. Unfortunately, we do not have an infinite number of corresponding physical principles needed to avoid the problem.

Therefore, one must either indicate new (sufficiently powerful) principles or radically reduce the number of free parameters in the theory. Anyway, we need to know the whole list of free parameters which S-matrix depends upon.

- Technically speaking, we need to make all Green functions UV finite, including $\dim = 6$ operators and, at least, $\dim = 8$ operators. This means computing the UV divergent part of a huge number of diagrams. The problem is already clear for the $\dim = 4$ part of the Lagrangian. To give an example, think about one-loop CTs for QGR. Is there a simple path?

Yes, it was *invented* by G. 't Hooft and it is based on the so-called background-field-method (BFM).

- For the SMEFT the problem is even more severe, since the number of CTs grows with the order in PT. Any calculation will produce a huge number of terms, most of them being related by EoMs or differing by a total divergence. Consider the following Lagrangian

Consider the following Lagrangian

$$\begin{aligned}\mathcal{L}(A + \phi) &= \mathcal{L}(A) + \phi_i \mathcal{L}'_i(A) + \frac{1}{2} \partial_\mu \phi_i W_{ij}^{\mu\nu}(A) \partial_\nu \phi_j \\ &+ \phi_i N_{ij}^\mu \partial_\mu \phi_j + \frac{1}{2} \phi_i M_{ij}(A) \phi_j + \mathcal{O}(\phi^3) + \text{total derivative}\end{aligned}$$

Classical equations of motion are $\mathcal{L}'_i(A) = 0$. All one loop diagrams are generated by $\mathcal{L}(\phi)$, the part quadratic in ϕ .

Assume

$$W_{ij}^{\mu\nu} = -\delta^{\mu\nu} \delta_{ij}$$

$$\mathcal{L}(\phi) = -\frac{1}{2} (\partial_\mu \phi)^2 + \phi N^\mu \partial_\mu \phi + \frac{1}{2} \phi M \phi$$

The Counter- Lagrangian is

$$\begin{aligned} \Delta \mathcal{L} = & \frac{1}{8\pi^2(n-4)} \left[a_0 M^2 + a_1 (\partial_\mu N_\nu)^2 + a_2 (\partial_\mu N_\mu)^2 + a_3 M N^2 \right. \\ & \left. + a_4 N_\mu N_\nu \partial_\mu N_\nu + a_5 (N^2)^2 + a_6 (N_\mu N_\nu)^2 \right] \end{aligned}$$

However

$$\mathcal{L}(\phi) = -\frac{1}{2} (\partial_\mu \phi + N_\mu \phi)^2 + \frac{1}{2} \phi X \phi$$

$$X = M - N^\mu N_\mu$$

is invariant under

$$\begin{aligned}\phi' &= \phi + \Lambda \phi \\ N'_\mu &= N_\mu - \partial_\mu \Lambda + [\Lambda, N_\mu] \\ X' &= X + [\Lambda, X]\end{aligned}$$

Therefore $\Delta \mathcal{L}$ also will be invariant

$$\Delta \mathcal{L} = \frac{1}{\varepsilon} \text{Tr} (a X^2 + b Y^{\mu\nu} Y_{\mu\nu})$$

$$Y_{\mu\nu} = \partial_\mu N_\nu - \partial_\nu N_\mu + N_\mu N_\nu - N_\nu N_\mu$$

Only two invariant scalars

To understand the EFT problem we consider a toy model, one complex scalar field, but will include both $\text{dim} = 6$ and $\text{dim} = 8$ operators. The Lagrangian looks as follows:

$$\mathcal{L} = \mathcal{L}_4 + \frac{g_6}{M^2} \mathcal{L}_6 + \frac{g_8^2}{M^4} \mathcal{L}_8$$

$$\mathcal{L}_4 = -\partial_\mu \phi^* \partial_\mu \phi - m^2 \phi^* \phi - \frac{1}{2} \lambda (\phi^* \phi)^2$$

$$\begin{aligned} \mathcal{L}_6 = & g^2 \left[a_1 |\phi|^2 \square |\phi|^2 + a_2 (\phi^* \partial_\mu \phi)^* \phi^* \partial_\mu \phi \right. \\ & \left. + g^2 a_3 \left(|\phi|^2 - \frac{1}{2} v^2 \right)^3 \right] \end{aligned}$$

$$\begin{aligned}
\mathcal{L}_8 &= g^6 a_4 \left(|\phi|^2 - \frac{1}{2} v^2 \right)^4 + g^4 a_5 \left(|\phi|^2 - \frac{1}{2} v^2 \right)^2 B \\
&+ g^4 a_6 |\phi|^2 V_\mu^* V_\mu + g^2 a_7 B^2 \\
&+ \frac{1}{2} g^2 a_8 \left(V_\mu^* U_\mu + V_\mu U_\mu^* \right) + a_9 S B \\
&+ a_{10} T_{\mu\nu} T_{\nu\mu} + a_{11} Z_{\mu\nu}^* Z_{\mu\nu} + a_{12} (\text{Re} Z_{\mu\nu}) (\text{Re} T_{\mu\nu})
\end{aligned}$$

$$B = 2 \partial_\mu \phi^* \partial_\mu \phi + \phi^* \square \phi + \phi \square \phi^*$$

$$V_\mu = \phi^* \partial_\mu \phi$$

$$U_\mu = (\square \phi^*) \partial_\mu \phi + 2 \partial_\nu \phi^* \partial_\mu \partial_\nu \phi + \phi^* \square \partial_\mu \phi$$

$$S = \partial_\mu \phi^* \partial_\mu \phi$$

$$T_{\mu\nu} = \partial_\mu \phi^* \partial_\nu \phi$$

$$Z_{\mu\nu} = \phi^* \partial_\mu \partial_\nu \phi$$

Introduce

$$\begin{aligned}\phi &= \frac{1}{\sqrt{2}} (h + v + i\psi) \\ m^2 &= \beta_h - \frac{1}{2}v^2 \\ v &= \frac{M}{g}\end{aligned}$$

where β_h fixes tadpoles

However, the Lagrangian is not canonically normalized (LSZ etc.), therefore we have to transform the fields, which is allowed by the Equivalence Theorem (ET)

$$h = \left(1 + g_6 x_h^{(1)} + g_6^2 x_h^{(2)}\right) h' + \frac{g_6^2}{M^2} X_h \square h'$$

$$\psi = \left(1 + g_6 x_\psi^{(1)} + g_6^2 x_\psi^{(2)}\right) \psi' + \frac{g_6^2}{M^2} X_\psi \square \psi'$$

$$M = \left(1 + g_6 x_M^{(1)} + g_6^2 x_M^{(2)}\right) M'$$

$$x_M^{(1)} = -x_h^{(1)} \quad x_h^{(1)} = \frac{1}{4} a_2 - a_1 \quad x_\psi^{(1)} = \frac{1}{4} a_2$$

$$X_\psi = \frac{1}{4} (a_8 - a_{11}) \quad x_M^{(2)} = -x_h^{(2)} + a_1^2 - \frac{1}{2} a_1 a_2 + \frac{1}{16} a_2^2$$

$$X_h = -a_7 + \frac{1}{4} a_8 - \frac{1}{4} a_{11} \quad x_\psi^{(2)} = \frac{3}{32} a_2^2 + \frac{1}{8} a_6$$

$$x_h^{(2)} = \frac{3}{2} a_1^2 - \frac{3}{4} a_1 a_2 + \frac{3}{32} a_2^2 + \frac{1}{8} a_6 - a_7 + \frac{1}{4} a_8 - \frac{1}{4} a_{11}$$

BFM obtains by expanding, $h \rightarrow H + h, \psi \rightarrow \Psi + \psi$. The resulting Lagrangian is still invariant under the following (shifted) transformations

$$H \rightarrow H + \Lambda \Delta H \quad \Psi \rightarrow \Psi + \Lambda \Delta \Psi$$

$$\begin{aligned} \Delta H &= -\Psi - g_6 a_1 \Psi \\ &+ g_6^2 \left[\left(\frac{1}{2} a_1^2 - \frac{1}{2} a_1 a_2 - a_7 + \frac{1}{4} a_8 - \frac{1}{4} a_{11} \right) \Psi - \frac{a_7}{M^2} \square \Psi \right] \\ \Delta \Psi &= H + \frac{M}{g} + g_6 \left[\frac{1}{2} (2 a_1 - a_2) \frac{M}{g} - a_1 H \right] \\ &+ g_6^2 \left[\frac{1}{4} (6 a_1^2 - 2 a_1 a_2 - 4 a_7 + a_8 - a_{11}) H \right. \\ &\left. - \frac{1}{4} (2 a_1^2 + a_6 - 4 a_7 + a_8 - a_{11}) \frac{M}{g} - \frac{a_7}{M^2} \square H \right] \end{aligned}$$

Remarks:

- Once we have the Lagrangian written in the language of the BFM we can use EoMs and the relevant message is that EoMs must only be used for classical fields
- Next, the integration over the quantum fields is Gaussian and can be performed according to a well-known algorithm
- It is worth noting that the problem of finding UV poles is reduced to the one of computing tadpoles
- Still we will have plenty of terms that are equivalent when using EoMs and neglecting total derivatives. We need an algorithm for that
- Only at this point we can determine field/parameter CTs and the mixing among Wilson coefficients. The presence of $\dim = 8$ operators is not trivial

Redundancy algorithm

$$\text{TD}_1 = \partial_\mu \partial_\mu \partial_\nu \partial_\nu H^2 = 0$$

...

$$\text{TD}_{18} = \partial_\mu (\partial_\mu \partial_\nu \Psi) \partial_\nu \Psi = 0$$

Apply EoMs and introduce scalars

$$B_1 = (\partial_\mu \partial_\nu H)^2$$

$$B_2 = (\partial_\mu \partial_\nu \Psi)^2$$

$$B_3 = (\partial_\mu \partial_\nu H) (\partial_\mu \partial_\nu \Psi)$$

$$B_4 = (\partial_\mu H) (\partial_\mu \Psi)$$

Solve and obtain

$$B_1 = -M^2 K_h + \mathcal{O}(\text{fields}^3) + \mathcal{O}(\text{fields}^4)$$

etc.

$$K_h = \partial_\mu H \partial_\mu H$$

Move to 3 fields, e.g.

$$\text{TD}_1 = \partial_\mu \partial_\mu \partial_\nu \partial_\nu H^3 = 0$$

...

$$\text{TD}_{46} = \partial_\mu (\partial_\mu \Psi) H^2 = 0$$

Apply EoMs, introduce scalars

$$C_1 = H (\partial_\mu \partial_\nu H)^2$$

...

$$C_{20} = \Psi \partial_\mu \Psi \partial_\mu \Psi$$

Solve the equations, e.g.

$$C_1 = \frac{1}{4} M^4 H^3 + \text{fields}^4 + \dots + \text{fields}^7$$

until all redundant expressions are eliminated.

The Lagrangian has the following structure

$$\begin{aligned}
 \mathcal{L} &= \sum_{n=0,2} g_6^n M_{0n}(1,1) h^2 + \sum_{n=0,2} g_6^n M_{0n}(2,2) \psi^2 \\
 &+ \sum_{n=0,2} g_6^n M_{0n}(1,2) h \psi + \sum_{n=1,2} g_6^n M_{1n}^\mu(1,2) h \partial_\mu \psi \\
 &\dots \\
 &+ g_6^2 M_4(1,2) h \square \square \psi
 \end{aligned}$$

Gaussian integration amounts to compute the trace of a logarithm where, in the M-matrices one replaces

$$\partial_\mu \rightarrow \partial_\mu + i q_\mu$$

integration over q follows

Remark: we are looking for UV divergent parts, therefore the expansion stops after a finite number of terms. Typical integrals (tadpoles) are:

$$\int d^n q \frac{1}{(q^2 + M^2)^j} = F(j; M)$$

...

$$\int d^n q \frac{q^{\mu_1} \dots q^{\mu_8}}{(q^2 + M^2)^j} = J(j; M) \delta^{\mu_1 \dots \mu_8}$$

Only $j = 1, 2$ give UV poles, etc.

$$\begin{aligned}
\ln M(x, \partial_x) \delta(x-y) &= \ln M(x, \partial_x) \int \frac{d^4 q}{(2\pi)^4} \exp\{i q \cdot (x-y)\} \\
&= \int \frac{d^4 q}{(2\pi)^4} \exp\{i q \cdot (x-y)\} \ln M(x, \partial_x + i q)
\end{aligned}$$

we have to compute

$$\exp\left\{-\frac{1}{2} \text{Tr} \ln M \delta(x-y)\right\}$$

and we perform an expansion of the trace in inverse powers of propagators. Since we are looking for UV poles, the expansion will stop somewhere (up to M^6 is required when $\text{dim} = 8$ operators are inserted).

It is worth noting that $\text{Tr} \ln M$ requires a symmetric matrix M and that particular care is due in avoiding spurious IR divergences (ψ is massless)

$$\ln(AB) = \ln A + \ln B \quad \text{if} \quad [A, B] = 0$$

$$\text{Tr} \ln(AB) = \text{Tr} \ln A + \text{Tr} \ln B \quad \text{if} \quad AB \text{ positive-definite}$$

Solution: write $M = \lambda (I + K)$ and use

$$\ln[\lambda (I + K)] = \ln(\lambda I) + \ln(I + K) = (\ln \lambda) I + K - \frac{1}{2} K^2 + \dots$$

Only at this point and after eliminating all redundant terms, we substitute (CTs)

$$\begin{aligned} H &= Z_h H & \Psi &= Z_\psi \Psi \\ M &= Z_M M & g &= Z_g g \end{aligned}$$

$$\begin{aligned} Z_h &= 1 + g^2 \left(\delta Z_h^{(4)} + g_6 \delta Z_h^{(6)} + g_6^2 \delta Z_h^{(8)} \right) \frac{1}{\epsilon} \\ &\dots \\ Z_g &= 1 + g^2 \left(\delta Z_g^{(4)} + g_6 \delta Z_g^{(6)} + g_6^2 \delta Z_g^{(8)} \right) \frac{1}{\epsilon} \end{aligned}$$

$$\delta Z_g^{(6)} = \sum_{n=1,3} \delta Z_g^{(6)}(n) a_n^r$$

$$\begin{aligned} \delta Z_g^{(8)} &= \delta Z_g^{(8)}(1,1) a_1^r a_1^r + \delta Z_g^{(8)}(2,2) a_2^r a_2^r + \delta Z_g^{(8)}(1,2) a_1^r a_2^r \\ &+ \delta Z_g^{(8)}(3,3) a_3^r a_3^r + \delta Z_g^{(8)}(1,3) a_1^r a_3^r + \delta Z_g^{(8)}(2,3) a_2^r a_3^r \\ &+ \sum_{j=4,12} \delta Z_g^{(8)}(j) a_j^r \end{aligned}$$

and mixings

$$\begin{aligned} a_i &= \sum_{j=1,3} \mathcal{L}_{ij} a_j^r + \sum_{j=4,12} \mathcal{L}_{ij}^{(6)} a_j^r \\ &+ \sum_{j=1,3} \sum_{l=j,3} \mathcal{L}_{ijl}^{(6)} a_j^r a_l^r \quad i = 1, 2, 3 \end{aligned}$$

$$a_i = \sum_{j=4,12} \mathcal{L}_{ij} a_j^r + \sum_{j=1,3} \sum_{l=j,3} \mathcal{L}_{ijl} a_j^r a_l^r$$

$$\mathcal{L}_{ij} = \delta_{ij} + g^2 \frac{1}{\varepsilon} C_{ij}$$

$$\mathcal{L}_{ijl} = g^2 \frac{1}{\varepsilon} C_{ijl}^{(6)}$$

$$\mathcal{L}_{ij}^{(6)} = g^2 g_6 \frac{1}{\varepsilon} C_{ij}^{(6)}$$

$$\mathcal{L}_{ijl}^{(6)} = g^2 g_6 \frac{1}{\varepsilon} C_{ijl}^{(6)}$$



Finally:

dim = 4

$$\delta Z_h^{(4)} = \delta Z_\psi^{(4)} = 0 \quad \delta Z_M^{(4)} = \delta Z_g^{(4)} = -10$$

dim = 6

$$\begin{aligned} \delta Z_h^{(6)}(i) &= 0 \\ \delta Z_\psi^{(6)}(1) &= 4 \quad \delta Z_\psi^{(6)}(2) = 2 \quad \delta Z_\psi^{(6)}(3) = 0 \\ \delta Z_M^{(6)}(1) &= 108 \quad \delta Z_M^{(6)}(2) = -26 \quad \delta Z_M^{(6)}(3) = -66 \\ \delta Z_g^{(6)}(1) &= 112 \quad \delta Z_g^{(6)}(2) = -24 \quad \delta Z_g^{(6)}(3) = 66 \\ C_{11} &= -4 \quad C_{12} = -2 \quad C_{13} = 0 \\ C_{21} &= -24 \quad C_{22} = -12 \quad C_{23} = 0 \\ C_{31} &= -104 \quad C_{32} = 24 \quad C_{33} = -76 \end{aligned}$$

A longer solution for dim = 8



Thank you for your attention

Backup Slides

EoMs: any change in the counter-Lagrangian which is proportional to \mathcal{L}' (classical fields) does not influence the S-matrix. In other words: one may make use of EoMs to simplify $\Delta \mathcal{L}$

$$\square H = H_4 + \mathcal{O}(g_6) \quad \square \Psi = P_4 + \mathcal{O}(g_6)$$

$$H_4 = M^2 H + \frac{3}{2} g M H^2 + \frac{1}{2} g^2 H^3 + \frac{1}{2} g^2 H \Psi^2 + \frac{1}{2} g M \Psi^2$$

$$P_4 = g M H \Psi + \frac{1}{2} g^2 H^2 \Psi^2 - \frac{1}{2} g^2 \Psi^3$$

$$Z = \int \mathcal{D}\phi \exp\{-\langle \phi, D\phi \rangle\} = \det^{-1/2}(D)$$

requires D to be self-adjoint

$$\langle \phi_1, D\phi_2 \rangle = \langle D\phi_1, \phi_2 \rangle$$

for any ϕ_1, ϕ_2