

# Future pp/hh and ep/h Colliders

Preface

HL LHC and its ep/A complement

FCC-hh with its associated eh option

HE LHC

Personal Remarks on the Physics, Accelerator and Projects

Max Klein (U Liverpool)  
ATLAS, H1, LHeC and FCC-eh

Invited Talk at the UK PPAP Meeting, Rutherford Laboratory, 20<sup>th</sup> of July, 2017

# 50 years ago



**Robert Jungk (1966)**

Die grosse Maschine  
auf dem Weg in eine andere Welt

**The big machine**


on the road into a new world

A book on the Proton Synchrotron ..



Niels Bohr at 1<sup>st</sup> Council 1952  
Council: highest level committee

ICHEP 1966



No Standard Model, Theory confused,  
ECFA, Amaldi: SPS for CERN  
Experiment paved the way:  
Quarks (ep)  $\rightarrow$  QCD,  $SU_L(2) \times U(1)$

# Funding HEP




Softened but hard  
Brexit from SPS in 68  
→ repeal shortly after

E.Amaldi to ECFA, 10.7.1968

In the Council meeting of 19 June, the United Kingdom delegation announced the decision of the British Government not to participate in the 300 GeV project. This decision was essentially based on economical considerations; the scientific and technical merits of the project were not questioned. The British delegate added a personal statement endorsed by the competent scientific authorities in his country in which as a physicist he regretted the decision of his Government and hoped that it would be possible at a later time to come back on it.

**convincing us, the academic and the public society – necessary, not always sufficient**

# The Current Major HEP Funding Resource is -CERN



CERN has  $O(200)$  MSF free money annually  $\rightarrow$  100 years to fund a 20 BSF project this way.



# Time Projections

## *Scientific activities*

European Strategy 2006

3. The LHC will be the energy frontier machine for the foreseeable future, maintaining European leadership in the field; *the highest priority is to fully exploit the physics potential of the LHC, resources for completion of the initial programme have to be secured such that machine and experiments can operate optimally at their design performance. A subsequent major luminosity upgrade (SLHC), motivated by physics results and operation experience, will be enabled by focussed R&D; to this end, R&D for machine and detectors has to be vigorously pursued now and centrally organized towards a luminosity upgrade by around 2015.*

Most likely, the LHC will have been the main base for HEP for ~50 years...

**Apparently we are unable to deliver reliable time projections  
... and yet we need optimism in order to progress ...**

## Particle Physics - a Sequence of Spectroscopies

- "Excitation of the 2536 Å Resonance Line of Mercury"  
Franck / Hertz 1914  
Bohr → ATOMIC SPECTROSCOPY
- "Disintegration of Elements by High Velocity Protons"  
Cockcroft / Walton 1932  
 $p\text{Li} \rightarrow \alpha\alpha$  : NUCLEAR SPECTROSCOPY
- "Total Cross-Sections of Positive Pions in Hydrogen"  
Anderson/Fermi/Long/Nagle 1952  
 $\Delta^{++} \rightarrow p\pi$  : HADRON SPECTROSCOPY
- The charming "November Revolution"  
Ting et al., Richter et al. 11.11.1974  
 $\mathcal{J}/\Psi \rightarrow c\bar{c}$ : QUARK SPECTROSCOPY



Gustav Hertz: Nobel 1925



John Cockcroft and Ernest Walton: Nobel 1951



Enrico Fermi: Nobel 1935



Sam Ting and Burt Richter: Nobel 1976

**No new spectroscopy appeared – neither 1992 (LEP) nor 2012 (LHC),  
No SUSY, neither at 100 GeV nor at 1000 GeV → a major surprise**

# The Standard Model is (formally) complete

**SM was completed with a series of pp, ee and ep machines** exploring the 10 GeV scale (ISR, SppS – PETRA, Tristan – electron, muon and neutrino experiments) and the Fermi scale (Tevatron – LEP, SLC – HERA), **besides further dedicated experiments [ep SLAC78..]**.

All three types of colliding experiments were instrumental in the SM establishment:  
For example: LEP predicted the top mass and Tevatron found the top quark;  
HERA measured the gluon distribution and LHC discovered  $gg \rightarrow \text{Higgs} \rightarrow 4l, \gamma\gamma$ .  
Tevatron saw excess in high pt jets, yet attributed to PDFs with DIS etc

For the first time since decades we have NO definite guidance, no SM particle to find. Note, however, that the Tevatron, LEP and HERA proposals largely emphasised NOT the SM but the BSM (SUSY, LQ) physics. Rarely the SM was a funding argument before either and the theory was no less speculative. Theory only guides: e.g. Weinberg 1980 SU(5): end of colliders, go underground to see proton decay ... to find neutrino oscillations ..

The LHC stands alone, it has no ep partner to explore the 1 TeV scale and it has no ee partner to study the Higgs boson. Can we build in time a 1 TeV ep collider (yes we could) and can we build a higher (than LEP) energy ee collider (for Phil to discuss)

**The FCC study has hh, ee and eh: yet 5?: time, cost, technology, theory, detectors**  
+ the public acceptance of such a major step into the unknown and below Lac Lemman

# HL LHC



Machine: 950 MSF [170 CE, 120 in kind], LIU 180 MSF; 1800 CERN person years + 1100 FTE  
ATLAS: major detector upgrade and tracker replacement; CMS: major upgrades, fwd calo..

Total cost may be close to 2 BSF and involves 10k physicists in experiment and accelerators

The first priority of the 2020 strategy update will no doubt be the HL LHC, for 2 decades hence






# LIU project

## Increase injector reliability and lifetime to cover HL-LHC run (until ~2040) closely related to consolidation program

- ⇒ Upgrade/replace ageing equipment (power supplies, magnets, RF...)
- ⇒ Improve radioprotection measures (shielding, ventilation...)

## Increase intensity/brightness in the injectors to match HL-LHC requirements

- ⇒ Enable Linac4/PSB/PS/SPS to accelerate and manipulate higher intensity beams (efficient production, space charge & electron cloud mitigation, impedance reduction, feedbacks, etc.)
- ⇒ Upgrade the injectors of the ion chain (Linac3, LEIR, PS, SPS) to produce beam parameters at the LHC injection that can meet the luminosity goal



	$\mathcal{N}$ ( $\times 10^{11}$ p/b)	$\epsilon$ ( $\mu\text{m}$ )
LIU Baseline	2.3	2.2
HL-LHC	2.3	2.1

# LHC Upgrade Goals: Performance optimization

Luminosity recipe (round beams):

$$L = \frac{n_b \cdot N_1 \cdot N_2 \cdot \gamma \cdot f_{rev}}{4\pi \cdot \beta^* \cdot \varepsilon_n} \cdot F(\phi, \beta^*, \varepsilon, \sigma_s)$$

- 1) maximize bunch intensities
- 2) minimize the beam emittance
- 3) minimize beam size (constant beam power); → triplet aperture
- 4) maximize number of bunches (beam power); → 25ns
- 5) compensate for 'F'; → Crab Cavities
- 6) Improve machine 'Efficiency' → minimize number of unscheduled beam aborts


→ Injector complex

Upgrade LIU

→ Crab Cavities

→ minimize number of unscheduled beam aborts

# The critical zones around IP1 and IP5



3. For collimation we also need to change the DS in the continuous cryostat in IR7:  
11T Nb<sub>3</sub>Sn dipole


2. We also need to modify a large part of the matching section  
e.g. Crab Cavities & D1, D2

1. New triplet Nb<sub>3</sub>Sn required due to:  
-Radiation damage  
-Need for more aperture


Changing the triplet region is not enough for reaching the HL-LHC goal!

→ More than 1.2 km of LHC !!  
→ Plus technical infrastructure (e.g. Cryo and Powering)!!

# IR1 & IR5 Underground Civil Engineering:



P. Fessia, HL-LHC TDR







today

# LHC / HL-LHC Plan




High  
Luminosity  
LHC



FAV = Fabrication, Assembly and Verification




# Collider Luminosities vs Year (pp and ep)



# Parameters of CERN pp Colliders

parameter	FCC-hh		HE-LHC	(HL) LHC
collision energy cms [TeV]	100		27	14
dipole field [T]	16		16	8.33
circumference [km]	100		27	27
straight section length [m]	1400		528	528
# IP	2 main & 2		2 & 2	2 & 2
beam current [A]	0.5		1.12	(1.12) 0.58
bunch intensity [ $10^{11}$ ]	1	1 (0.2)	2.2 (0.44)	(2.2) 1.15
bunch spacing [ns]	25	25 (5)	25 (5)	25
rms bunch length [cm]	7.55		7.55	(8.1) 7.55
peak luminosity [ $10^{34} \text{ cm}^{-2}\text{s}^{-1}$ ]	5	30	25	(5) 1
events/bunch crossing	170	1k (200)	~800 (160)	(135) 27
stored energy/beam [GJ]	8.4		1.3	(0.7) 0.36
beta* [m]	1.1-0.3		0.25	(0.20) 0.55
norm. emittance [ $\mu\text{m}$ ]	2.2 (0.4)		2.5 (0.5)	(2.5) 3.75

## HIGGS PHYSICS AT THE LHeC SUMMARY




- GLUON FUSION AND  $W$  FUSION  $\Rightarrow$  PDF+ $\alpha_s$  UNCERTAINTY REMOVED (hatched bands)
- $H\bar{b}b$  MEASURED TO PERCENTAGE PRECISION;
- $\tau\tau$  AND  $\bar{c}c$  ALSO MEASURABLE

S.Forte ECFA 11/15


The exp. error on the Higgs cross section calculated with LHeC PDF is 0.3%  $\rightarrow$  sensitive to mass

# Extended Search Programme (SUSY?)



5 $\sigma$  up to  $\sim 2.5 \text{ TeV}$  gluinos  
@ HL-LHC


ATLAS-PUB-2014-10



5 $\sigma$  up to  $\sim 3 \text{ TeV}$  squarks  
5 $\sigma$  up to  $\sim 1.2 \text{ TeV}$  stops  
5 $\sigma$  up to  $\sim 1.3 \text{ TeV}$  sbottoms  
@ HL-LHC

SUSY is too beautiful to not exist but it is broken “heavier and heavier”

Physics studies of the HL LHC Potential: ongoing  $\rightarrow$  **HL/HE LHC Workshop 17/18**



# LHC Full Energy Exploitation:

Operation beyond ultimate beam energy:  $E > 7.56\text{TeV}$

Proposal to replace 1/3 of all MB magnets with 11T Nb<sub>3</sub>Sn magnets

- Major interventions [opening of all MB interconnects]
- Not clear other magnets can be scaled up in energy  
e.g. insertion quadrupoles and triplet magnets
- Not clear other systems [e.g. beam dump system] can be easily upgraded


Study ongoing and results expected for end 2018 / beginning 2019

Could perhaps be envisaged as a second LHC upgrade at the end of the HL-LHC exploitation period. But investment [time and capital] will be substantial



# Add an Electron Beam (ERL) to LHC (+FCC)



Conceptual Design Report (2012), Update for next European Strategy



**Concurrent operation to pp, LHC/FCC become 3 beam facilities.  $P(e) < 100$  MW  
 $10^{34}$  luminosity and factor of 20 (LHC) extension of  $Q^2$ ,  $1/x$  reach vs HERA**



# Optimize Cost vs Physics and Time



A rough extrapolation of a 3-turn ERL shows how the cost rises non-linearly with the electron beam energy. Reliable cost estimate work in progress

9km: 1/3 of U(LHC) leads to 60 GeV e energy  
 5.4km : 1/5 of LHC circumference: 51 GeV

**Conclusion on LHeC: may build an ERL tangential to LHC (HL and HE in sight).  
 Choice of energy from optimization of physics, cost, effort, time schedule..**

# Five Major Themes of electron-hadron Physics

Cleanest High Resolution Microscopes

Joint ep and pp Physics

High Precision Higgs Exploration

Discovery Beyond the Standard Model


A Unique Nuclear Physics Facility

# Empowering pp Discoveries

External, reliable input (PDFs, factorisation..) is crucial for range extension + CI interpretation



GLUON

SUSY, RPC, RPV, LQS..




QUARKS

Exotic+ Extra boson searches at high mass



# Higgs Physics with ep



High cross section (cc: LHeC 200fb, FCC-eh 1pb)

Electroweak production, uniquely CC vs NC

Access to WW-H-WW and ZZ-H-ZZ

No pileup, clean theory, challenging simulations

## SM coupling measurement expectations

$\kappa$ in %	HL LHC	LHeC HL	LHeC HE	FCC-eh
$H \rightarrow bb$	10?	0.5	0.3	0.2
$H \rightarrow cc$	50??	4	2.8	1.8

Expected number of signal events  
( $E_e = 60$  GeV)

FCC ep (~85,000  $H \rightarrow bb$  events)

DLHC (~35,000  $H \rightarrow bb$  events)

LHeC (~15,000  $H \rightarrow bb$  events)


Recent Higgs-in-ep studies for CDR: Higgs self coupling from FCC-eh associated top-Higgs production, Higgs into invisible (dark matter), Exotic Higgs physics: H into light scalars,  $H^-$  and others

cf U Klein at FCC Berlin for references and summary

ep when added to pp turns the pp colliders into high precision Higgs facilities.  
Removes PDF and coupling constant uncertainties in pp gg fusion process.

# Possible Discoveries Beyond SM with eh

Search for Sterile Neutrinos  
(LHC/FCChh FCCee LHeC/FCCeh)



**QCD:**

(No) saturation of the gluon density

QCD radiation pattern (BFKL?) – hh!

New QCD states (instantons)

Higher symmetry embedding QCD

**Electroweak:**

EFTs, CI to 300 TeV, RPV SUSY

Exotic Higgs Decays (Dark Matter..)

Extension of Higgs Sector ( $H^{++}$ ..)


Sterile Neutrinos ...

# Powerful ERL for Experiments (ep, $\gamma\gamma$ ): PERLE at Orsay


PERLE at Orsay (LAL/INP) Collaboration: BINP, CERN, Daresbury/Liverpool, Jlab, Orsay +

3 turns, 2 Linacs, 400 MeV, 15mA, 802 MHz, Energy Recovery Linac facility

- Demonstrator of ERL for ep at LHC/FCC
- SCRF Beam based development facility
- Low E electron and photon beam physics
- High intensity:  $O(100) \times \text{ELI}$



CDR to appear in J Phys G [arXiv:1705.08783]




A.Bogacz

**Strong low energy physics program:**

p radius,  $\sin^2\theta$ , dark photons, photon-nuclear physics, ..

# FCC







# Implementation - new footprint baseline

Alignment Shafts Query

Choose alignment option  
V4variation\_v2017-2

Tunnel elevation at centre: 322mASL


Grad. Params

Azimuth (°): -23.5  
Slope Angle x-x(%): 0.3  
Slope Angle y-y(%): 0.08

LOAD SAVE CALCULATE

Alignment centre  
X: 2499941 Y: 1107760

	Angle	Depth	Angle	Depth
LHC	37°	49m	40°	83m
SPS		121m		126m
T12		121m		126m
T18		51m		118m




Geology intersected by Shafts				Shaft Depths		
Point	Actual	Shaft Depth (m)			Geology (m)	
		Molasse SA	Wildflysch	Quaternary	Molasse	Urgonian
A	152	0	0	0	152	0
B	121	0	0	26	95	0
C	127	0	0	44	83	0
D	205	66	0	40	100	0
E	89	0	0	89	0	0
F	476	0	0	49	427	0
G	307	0	0	73	234	0
H	266	0	0	0	266	0
I	198	0	0	11	187	0
J	248	0	0	1	247	0
K	88	0	0	70	18	0
L	172	0	0	89	83	0
Total	2449	66	0	492	1892	0

Optimisation in view of accessibility surface points, tunneling rock type, shaft depth, etc.

## Tunneling

- Molasse 90%, Limestone 5%, Moraines 5%



## Shallow implementation

- ~ 30 m below lakebed
- Reduction of shaft length and technical installations
- One very deep shaft F (RF or collimation), alternatives being studied, e.g. inclined access




# Designing the FCC-hh

- **synchrotron radiation power** → associated cryogenics power (after various technological improvements and mitigation – higher beam-screen temperature etc.) **limits** maximum number of protons (e.g.  $10^{15}$  p/beam or beam current of  $\sim 0.5$  A for 100 km ring at 100 TeV c.m.)
- maximum **beam current** + “turnaround time” (FCC + inj.) **constrain** integrated luminosity
- maximum **peak pile up**  $\sim 1000$  or  $\sim 200$  **limits** peak luminosity at 25 ns and 5 ns **bunch spacing**, respectively
- maximum **acceptable beam-beam tune shift** 0.01 or 0.03 + **optics** (minimum  $\beta^* \sim 0.3$ -1.1 m) **also limits** peak luminosity

# The potential of a Future Circular Collider



- Guaranteed deliverables:
  - study of Higgs and top quark properties, and exploration of EWSB phenomena, with unmatched precision and sensitivity
- Exploration potential:
  - mass reach enhanced by factor  $\sim E / 14 \text{ TeV}$  (will be 5–7 at 100 TeV, depending on integrated luminosity)
    - *statistics enhanced by several orders of magnitude for BSM phenomena brought to light by the LHC*
  - benefit from both direct (large  $Q^2$ ) and indirect (precision) probes
- Provide firm Yes/No answers to questions like:
  - is the SM dynamics all there is at the TeV scale?
  - is there a TeV-scale solution to the hierarchy problem?
  - is DM a thermal WIMP?
  - did baryogenesis take place during the EW phase transition?

# Theory to pave new ways



# Direct Discovery Potential of the FCC (pp)

## Eg SUSY and DM reach at 100 TeV




### First preliminary findings:

- possibility to find (or rule out) thermal WIMP DM candidates
- possibility to support (or exclude) EW baryogenesis

$$M_{\text{WIMP}} \leq 1.8 \text{ TeV} \left( \frac{g^2}{0.3} \right)$$

# Searching for no-lose items..

Strong 1<sup>st</sup> order phase transition required to generate and sustain the out of equilibrium generation of a baryon asymmetry during EW symmetry breaking




Strong 1<sup>st</sup> order phase transition  $\Rightarrow \langle \Phi_c \rangle > T_c$

In the SM this requires  $m_H \lesssim 80$  GeV.

Since  $m_H = 125$  GeV, **new physics**, coupling to the Higgs and effective at **scales  $O(\text{TeV})$** , must modify the Higgs potential to make this possible

# FCC-hh Detector Study




70m cavern  
18m shaft  
“only 1 BSF”


4T 10m solenoid  
Forward solenoids  
Silicon tracker  
Barrel ECAL Lar  
Barrel HCAL Fe/Sci  
Endcap HCAL/ECAL Lar  
Forward HCAL/ECAL Lar

W Riegler et al  
June 2017

CMS




FCC-hh






# Underground structures



First basic + very preliminary design using new dimensions from MDI WG (w/o detailed radioprotection treatment, access optimisation, detailed design over several floors)



# Draft Schedule Considerations





# FCC-ee RF staging scenario


"Ampere-class" machine

	V tot (GV)	n bunch	I beam (mA)
Z	0.2	91500	1450
W	0.8	5260	152
H	3	780	30
t	10	81	6.6

"high gradient" machine

Three sets of RF cavities to cover all options FCCee & Booster:

- Installation sequence comparable to LEP ( $\approx 30$  CM/shutdown)
- high intensity (Z, FCC-hh): 400 MHz mono-cell cav,  $\approx 1$  MW source
- high energy (H, t): 400 MHz four-cell cavities, also for W machine
- booster and t machine complement: 800 MHz four-cell cavities
- Adaptable 100MW, 400MHz RF power distribution system



FCC-ee count on seminal luminosity for el.weak and only later H measurements  
FCC-ee as "a first step" would defer the hadron-hadron collider to in +50 years..  
If we were now at Lausanne 1984, the LHC started operating 2009: +25 years



# Conceptual Design Report

## 1 – PHYSICS

Physics opportunities across all scenarios

## 2 Hadron Collider Summary

## 4 Lepton Collider Summary

## 6 High Energy LHC Summary

## 3 – Hadron Collider Comprehensive

Accelerator

Injectors

Technologies

Infrastructure

Operation

Experiment

eh

## 5 – Lepton Collider Comprehensive

Accelerator

Injectors

Technologies

Infrastructure

Operation

Experiment

## 7 – High Energy LHC Comprehensive

Accelerator

Injectors

Infrastructure

*Refs to FCC-hh, HL-LHC, LHeC*

- Required for end 2018, as input for European Strategy Update
- Common physics summary volume
- Three detailed volumes FCChh, FCCee, HE-LHC
- Three summary volumes FCChh, FCCee, HE-LHC





# he civil engineering studies : preferred location

## Why is experimental point L preferred?

### Positives:

- Low geological risk compared to other locations, anticipated tunnelling in molasse only.
- Close to current CERN site.
- FCC ring relatively shallow at this point, therefore shallower shafts.


### Remaining problems:

- Potential clash with injection lines needs to be studied.
- Located inside the FCC ring so integration with other structures to be studied.
- Depth below Rhone to be evaluated.



Civil engineering optimisation for FCC-he (J. L. Stanyard)








# FCC-he Detector Layout - Scaled Version of LHeC Detector

## Inner Dipoles and Solenoid



# HE LHC

many aspects extrapolated/copied from HL-LHC or FCC-hh

exceptions:

tunnel integration and magnet technology

- push for **compact 16 T** magnets (magnetic cryostat, shielding)
- HE-LHC ***Nb<sub>3</sub>Sn*** magnets must be bent - 5 mm horizontal orbit shift over 14 m

arc optics

- higher dipole filling factor to reach energy target → different arc optics
- relaxed strength of quadrupoles and sextupoles → different arc optics

straights

- low-beta insertions, longer triplet than HL-LHC,  $\beta^*$  reach
- collimation straights, FCC-hh scaling not applicable,  
warm dipole length increases w.r.t. to LHC; new approach needed?!
- extraction straights – length of kicker & septum sections

injector

- determined by extraction system, physical & dynamic aperture, impedance...




# Parameters of CERN pp Colliders

parameter	FCC-hh		HE-LHC	(HL) LHC
collision energy cms [TeV]	100		27	14
dipole field [T]	16		16	8.33
circumference [km]	100		27	27
straight section length [m]	1400		528	528
# IP	2 main & 2		2 & 2	2 & 2
beam current [A]	0.5		1.12	(1.12) 0.58
bunch intensity [ $10^{11}$ ]	1	1 (0.2)	2.2 (0.44)	(2.2) 1.15
bunch spacing [ns]	25	25 (5)	25 (5)	25
rms bunch length [cm]	7.55		7.55	(8.1) 7.55
peak luminosity [ $10^{34} \text{ cm}^{-2}\text{s}^{-1}$ ]	5	30	25	(5) 1
events/bunch crossing	170	1k (200)	~800 (160)	(135) 27
stored energy/beam [GJ]	8.4		1.3	(0.7) 0.36
beta* [m]	1.1-0.3		0.25	(0.20) 0.55
norm. emittance [ $\mu\text{m}$ ]	2.2 (0.4)		2.5 (0.5)	(2.5) 3.75

## FCC-hh

## HE-LHC


V. Mertens et al.



**6 m inner tunnel diameter**

main space allocation:

- 1200 mm cryo distribution line (QRL)
- 1500 mm installed cryomagnet
- 1600 mm cryomagnet magnet transport
- >700 mm free passage.



**3.8 m inner tunnel diameter**

main space allocation:


- 850 mm cryo distribution line (QRL)
  - 1200 mm installed cryomagnet
  - 1200 mm cryomagnet magnet transport
- challenging*

# Nb<sub>3</sub>Sn Magnet Development – 16 T

Cryogenic aperture requirements in magnet cold-mass for half-cell cooling (~105 m)

4 cool-down & warm-up tubes (DN30, PN50)

Bayonet heat exchanger (DN95)



- How close is the best of today to the FCC specification (irrespective of  $d_{\text{effective}}$  or RRR)?
- **The best is actually not too far away: 1675 A/mm<sup>2</sup> versus 1850 A/mm<sup>2</sup>**

Also strong SC RF developments and progress (cf E Jensen Berlin)

- The present cost is > 20 Euro/kAm @ 4.2 K, 16 T (FCC target is 5 Euro/kAm)
- Large production (~7-10 kt for FCC-hh, 2-3 kt for HE-LHC)
  - ITER : ~500 tons in total, 100 tons/year, 8 companies
  - HL-LHC : ~ 20 tons in total, initially 2 companies, now 1 company)

## Development

Collaboration agreement with **KEK- Japan**. Development of Nb<sub>3</sub>Sn wire at **Jastec** and **Furukawa**

Collaboration agreement with **KAT – Korea**. Development of Nb<sub>3</sub>Sn wire.

Collaboration agreement with **Bochvar Inst. – Russia**. Development of Nb<sub>3</sub>Sn wire at **TVEL**

# HE LHC Time Schedule

## Remark

We recall that the US Nb<sub>3</sub>Sn conductor program started in 1999 aiming at the same target cost as the one set for the FCC. After 5 years the program stopped: the target cost was not achieved, remaining about three times higher than the target (presently we are exactly in the same situation, with no advancement with respect to the outcome of that US program). However the program was very successful and resulted in practically doubling the critical current and decreasing the cost of the conductor by more than a factor of 2.

This is to say that a vigorous R&D program, probably over 5-10 years, will be necessary before a massive production for a HE-LHC or a FCC can start.

D Tommassini (June 17)

Production of HE LHC Components: O(10) years.

Injector: Currently scSPS disfavoured and SPS 450 GeV considered. 80 years old by 2050


Detectors: ATLAS and CMS at twice the proton beam energy – major upgrades to study

HL LHC: 3ab<sup>-1</sup> estimated duration: until 2038 – 2040

Dismantling LHC, Installing HE LHC O(10) years: **HE LHC in 2050** maybe a bit earlier.

**Total cost O(5) BSF:** 25 years of 200 MSF. Magnet cost crucial to reduce. **Physics ?**

# Projected Timelines for Future ep/eA Colliders



**HERA:** Proposal 1984, Data 1992-2007, Publications 1993-2018

**VHEep:** Plasma e – LHC. **Chinese ep/A** projects: Lanzhou (low E) and CEPC/SPPC

**Disclaimer:** For discussion and illustration at DIS17 only MK+RY, April 7<sup>th</sup>, 2017, DIS at Birmingham

# Luminosity for LHeC, HE-LHeC and FCC-ep

parameter [unit]	LHeC CDR	ep at HL-LHC	ep at HE-LHC	FCC-he
$E_p$ [TeV]	7	7	12.5	50
$E_e$ [GeV]	60	60	60	60
$\sqrt{s}$ [TeV]	1.3	1.3	1.7	3.5
bunch spacing [ns]	25	25	25	25
protons per bunch [ $10^{11}$ ]	1.7	2.2	2.5	1
$\gamma\epsilon_p$ [ $\mu\text{m}$ ]	3.7	2	2.5	2.2
electrons per bunch [ $10^9$ ]	1	2.3	3.0	3.0
electron current [mA]	6.4	15	20	20
IP beta function $\beta_p^*$ [cm]	10	7	10	15
hourglass factor $H_{geom}$	0.9	0.9	0.9	0.9
pinch factor $H_{b-b}$	1.3	1.3	1.3	1.3
proton filling $H_{coll}$	0.8	0.8	0.8	0.8
luminosity [ $10^{33}\text{cm}^{-2}\text{s}^{-1}$ ]	1	8	12	15

Oliver Brüning<sup>1</sup>, John Jowett<sup>1</sup>, Max Klein<sup>1,2</sup>,  
Dario Pellegrini<sup>1</sup>, Daniel Schulte<sup>1</sup>, Frank Zimmermann<sup>1</sup>

<sup>1</sup> CERN, <sup>2</sup> University of Liverpool

April 6<sup>th</sup>, 2017

**ep at CERN: 1000 times  
the luminosity of HERA  
< 100 MW wall plug**



# Summary

A summary is personal and not straightforward. CERN has major opportunities and skills to drive the pp energy frontier higher + BSM. This can be substantially enriched by adding an ERL electron beam, at relatively low cost, for concurrent pp+ep (as well as AA+eA) operation, with the LHC, the HE LHC and the FCC.

The FCC appears as a vision while HL LHC is the next program. Working on the vision is important (infrastructure, magnets, RF, detectors... ). We shall invest in theoretical physics for without convincing guidance (no loose no-lose..) one will not be able to attract the world's attention + means for making that vision real.

In between, the HE LHC offers a next major step, based at large on FCC technology. That step is neither easy nor cheap and will be real hardly before 2050 unless the HL LHC programme will be substantially shortened, with an overriding justification..

Many thanks to many colleagues: M Benedikt, S Bertolucci, O Bruening, G Dissertori, E Jensen, U Klein, M Mangano, V Radescu, D Schulte, H Schopper, A Stocchi, F Zimmermann, + many others, the LHeC + all FCC Coordination teams, my colleagues at Liverpool and on ATLAS – all who join, tolerate or/and support thinking about the future.

## Sources and Events 2017

FCC Physics Week January, FCC at Berlin May, IAC FCC Review June, LHeC/FCC workshop September 11-13 (CERN), HL/HE LHC Workshop November 2017 .. 2018 (CERN)

title

# T Han Aspen

\* With the Higgs discovery, the SM is healthier than ever,  
valid to **a scale up to  $\Lambda \sim ?$**

But the Higgs sector fine-tuned  **$\delta$**  :

\* VLHC will take the lead for searches:


**$\tilde{g}, \tilde{t}, \tilde{b}, \chi^{\pm,0}, \dots H^{\pm}, A^0; W^{\pm'}, Z'$  ...**

The  **$top, W, Z, H$**  may hold the key for  
discovery!

• Searching for **new physics** starts from

understanding **old physics in the new regime**:

- **$top, W, Z$**  may behave as partons to produce new heavy states;
- **$top, W, Z, H$**  may serve as new radiation sources;  
and may help reveal new heavy states.
- Thus, need precise understanding of the dynamics/kinematics



# Future SUSY

Assuming a massless LSP

Model	Limit [TeV]	Discovery Reach [TeV]	
	8 TeV 20 fb <sup>-1</sup>	14 TeV 3000 fb <sup>-1</sup>	100 TeV 3000 fb <sup>-1</sup>
$pp \rightarrow \widetilde{g}\widetilde{g} \rightarrow q\bar{q}\widetilde{\chi}_1^0 q\bar{q}\widetilde{\chi}_1^0$	1.4 (ATLAS)	2.3	<b>11</b>
$pp \rightarrow \widetilde{g}\widetilde{g} \rightarrow t\bar{t}\widetilde{\chi}_1^0 t\bar{t}\widetilde{\chi}_1^0$	1.4 (ATLAS)	2.0	<b>6.0</b>
$pp \rightarrow \widetilde{q}\widetilde{q}^* \rightarrow q\widetilde{\chi}_1^0 \bar{q}\widetilde{\chi}_1^0$	1.0 (CMS)	1.0	<b>7.8</b>
$pp \rightarrow \widetilde{t}\widetilde{t}^* \rightarrow t\widetilde{\chi}_1^0 \bar{t}\widetilde{\chi}_1^0$	0.7 (CMS)	1.2 <sup>a</sup>	<b>6.5</b>

<sup>a</sup>[ATLAS projection](#)

M. Hance Aspen 15

SUSY is too beautiful to not exist but it is broken heavier and heavier

**For the FCC to be built we need overriding reasons which the society can accept for the project to go ahead. Magnets and theory are the main challenges of the FCC.**

# Report of the SSC Collider Dipole Review Panel

June 1989

SSC-SR-1040

G. Voss

Deutsches Elektronen-Synchrotron, DESY  
Hamburg, Germany

and

T. Kirk

SSC Central Design Group\*  
c/o Lawrence Berkeley Laboratory  
Berkeley, CA

design. The evaluation was based upon information provided in the scheduled topic presentations, comments and discussion from various Magnet Program personnel, and a set of documents provided by the SSC Magnet Systems Division head: *SSC Magnet R&D Plan 1988*, edited by E. L. Goldwasser; *Development Status for SSC Magnets*, December 1988; *SSC Magnet R&D Plan Update*, January 1989; and the SSC Magnet Program presentations given at the DOE SSC Annual Review, 30 January 1989.



The program goal is to provide a mature design for a 17-m-long magnet that is capable of producing a uniform dipole field with an intensity of 6.6 T at a temperature of 4.35 K and which satisfies all system requirements but is not yet optimized for industrial production. Further



# Development of High Field Dipoles



## HTS for 20 T

6 T HTS (YBCO) insert for test  
in FReSCa2 (no bore)



J.M. Rey, F. Borgnolutti, CEA-Saclay

A 20 T HE-LHC dipole  
E. Todesco, L. Rossi (CERN)



Cost optimized, graded winding

Sc wire: higher current, higher field. Reduced losses. NbTi: 15T at 10K, Nb<sub>3</sub>Sn: 25T at 4K, HTS inserts YBCO  
Cost is a major factor: today: Nb<sub>3</sub>Sn is 5 times the NbTi cost and HTS is 10 times Nb<sub>3</sub>Sn (O.Bruening at KET 2/17)  
HE LHC needs about 4500 tons of Nb<sub>3</sub>Sn, ITER needs 500 t




# HL LHC – Replacement of Inner Quadrupoles


Inner triplet quadrupoles receive 25MGy of radiation from  $300\text{fb}^{-1}$  of pp at the LHC →  
Larger aperture, larger field to ensure high luminosity performance: 1-2 decades of design




LHC: **NbTi technology**  
LHC (USA & JP, 5-6 m)  
 $\varnothing 70\text{ mm}$ ,  $B_{\text{peak}} \sim 8\text{ T}$   
1992-2005



> 10 years  
development for  
NbTi magnets




**LARP**  
HL-LHC: **Nb<sub>3</sub>Sn Technology**  
LARP TQS & LQ (4m)  
 $\varnothing 90\text{ mm}$ ,  $B_{\text{peak}} \sim 11\text{ T}$   
2004-2010




Ca. 20 years for  
Nb<sub>3</sub>Sn magnet  
development!


LARP HQ  
 $\varnothing 120\text{ mm}$ ,  
 $B_{\text{peak}} \sim 12\text{ T}$   
2008-2014




**LARP**



LARP & CERN  
MQXF  
 $\varnothing 150\text{ mm}$ ,  
 $B_{\text{peak}} \sim 12.1\text{ T}$   
2013-2020




**LARP**




title

# The 10-100 GeV Energy Scale [1968-1986]



# The Fermi Scale [1985-2010]



# Three conditions for prosperity of HEP

## Staff

Positions for next generations: CERN staff + visitors: 1960 1166 → 1965 2530

A major new step will depend on how we can keep HEP attractive for life plans.

## Accelerators

**L.D. Landau:**

**Accelerators have the advantage to control the initial conditions**



Europe's big machines

???

LHC

LEP, HERA

SppS, PETRA

SPS

time

Accelerators need sites and major institutions. CERN should better have strong European partners DESY, Frascati, RAL, Saclay,.. and global challenges too.



Robert Jungk (1966)

Die grosse Maschine  
-auf dem Weg in eine andere Welt

The big machine  
-on the road into a new world





# 5. Big Questions

- Do we have too many particles? 12 leptons, 36 quarks, 12 mediators, 1 Higgs = 61
- Is there a further layer of structure (preons?)
- How can we unify the 3 + 1 interactions (SU(5) failed in 1980 but established neutrino physics)
- Why are leptons and quarks different?
- Can one restore the boson-fermion symmetry (SUSY since 1972)
- Why do we have 3 families?
- Neutrino puzzles: Majorana, sterile neutrinos – Oscillations (98), Pontecorvo (57)
- Is the proton stable?
- ...
- And: what is “behind” dark matter.. ? Not sure that is a particle physics question?

**New: We lost the SM guidance**

**Reminds on Kelvin, Planck ~1900**

**Note: 500 ATLAS papers**

# 1. The increase of energy and luminosity often led to discoveries

## Substructure discovered at Stanford

Hofstadter et al: 1957:  $ep \rightarrow ep$   $E_e=200$  MeV beam: proton finite radius of 1fm

Taylor et al: 1968:  $ep \rightarrow eX$   $E_e=1-20$  GeV beam (2 mile linac): partons at 0.1fm

## W,Z Bosons discovered at CERN

ISR in 1970,

SPS in 1974  $E_p=450$  GeV (fixed target lh, hh experiments, injector for LHC)

transformed to SppS Collider  $L=10^{30-31} \text{ cm}^{-2} \text{ s}^{-1}$  by van der Meer + Rubbia


UA1, UA2: first full acceptance pp detectors to catch  $W \rightarrow l\nu$  and MET

Partons came unexpected - despite the Quark Model  
W,Z were predicted in  $SU_L(2) \times U(1)$  electroweak theory

# Realization of the LHeC

LHC

Physics and cost will determine footprint



Pt 2

Preyessin Site

SPS

1/3 Pt8

MK 6/14

Pt 1

Pt 8

Meyrin Site

LHeC  
Civil Engineering  
Different Options  
Fraction 1/3-1/4-1/5  
Pt2 and Pt8  
J.OSBORNE/L.FAISANDEL GS-SE-DOP

# title






# Concluding Remarks

1. Electron-hadron scattering has **five big themes** (Microscope, Higgs, Joint to pp, BSM, Nuclei)  
→ It thus has a unique place in High Energy Physics (reaching beyond these themes too..)
2. **ep empowers pp**: searches and high precision (e.g. Higgs in pp+ep – an especially rich mix)  
→ ep and pp can operate concurrently: should be seen, studied and understood together
3. QCD deserves major new development through novel exp input (ep + pp) and theory  
→ **QCD may fail and lead BSM**: non-linear evolution, higher/grand symmetry, breaking of factorisation, valence components of heavy quarks, free colour, instantons, substructure..
4. ions: **eA at CERN is to revolutionise nuclear dynamics** and structure physics  
→ chromodynamic understanding of QGP, an EIC requires highest energy to be of highest value
5. Detector: one in LR, two in RR (HE-LHC?) **novel experimental opportunity post HL LHC upgrade**
6. **PERLE**: in time and scope to learn how to build and operate the ERL at high energy  
→ Electron-hadron configuration: genuine, high, added + crucial, **unique value for HEP**  
→ The ERL at the HL/HE/FCC is **affordable, i.e. it does NOT affect larger scale decisions** but possibly provides time until those may be taken. The electron energy is a function of available cost (in building and operating the ERL). The ep cms energy is much higher than that of ILC/CepC or FCCee (even CLIC in the FCC-eh case)  
→ **ep is an exciting, realistic option for a next energy frontier collider for particle physics**

# Intensity and Energy Frontier of Future DIS

## Lepton-Proton Scattering Facilities



From CERN Courier  
MK, H.Schopper  
June 2014


With input from  
A.Hutton, R.Ent,  
F.Maas, T.Rosner

**CERN: LHC+FCC: the only realistic opportunity for energy frontier deep inelastic scattering**  
**Huge step in energy ( $Q^2, 1/x$ ) and 2-3 orders of magnitude higher luminosity than HERA**




# High Precision for the LHC

## Higgs pp Cross Section



Predict the Higgs cross section in pp to 0.2% precision which matches the  $M_H$  measurement and removes the PDF error

ep+pp deliver high precision of Higgs and qcd and electroweak physics – compl to ee





D.Britzger on Thursday

Spacelike  $M_W$  to 10 MeV from ep  
→ Electroweak test at 0.01% !

Predict  $M_W$  in pp to 2.8 MeV →  
Remove PDF uncertainty on  $M_W$  in pp

Reach for  
 $\Lambda$  (CI ee qq):



LHeC: see CDR 2012

FCC - rough scaling only – very preliminary