

Higgs Properties

Victoria Martin
University of Edinburgh
UK ESU meeting Durham, April 2018



Disclaimers

- (Proto-)experiments have not investigated every their potential to measure every Higgs property. Many studies are still underway.
- LHC:
 - ➔ Generally results include an estimate of systematic effects
 - ➔ Most HL-LHC results presented are for one-experiment, expect $\sqrt{2}$ improvements for ATLAS+CMS.
 - ➔ ATLAS+CMS will extrapolate current Run 2 results for a Yellow Report at the end of this year
 - ➔ I didn't find any projections for HE-LHC
- Future experiments: ILC, CLIC, FCCxy
 - ➔ Most important is the projectiles, energy & luminosity
 - ➔ some are generator-level studies
 - ➔ systematic effects generally not estimated: statistical errors only
 - ➔ I've not looked for CEPC projections

What we want to know!

Is our 125 GeV Higgs boson the Standard Model Higgs boson?

If not, what is it?

Mass



Width

Spin / CP



Rares decay

Differential Cross Sections

Higgs boson cross sections: e^+e^-



Higgs boson cross sections: *pp*



Higgs boson mass

Compare Higgs boson mass to the couplings to W and Z :
check for consistency in the Standard Model

$$\delta(g_{HWW}) = 6.9\delta m_H \ ; \quad \delta(g_{HZZ}) = 7.7\delta m_H$$

Higgs boson mass @ lepton collider

ILC250: Reconstruct recoil mass from ZH

$$M_X^2 = (p_{CM} - (p_{\ell^+} + p_{\ell^-}))^2,$$



with 2 ab^{-1} : $m_H = \pm 14 \text{ MeV}$

CLIC: Measure Higgs mass directly from $H \rightarrow b\bar{b}$:


$$\Delta(m_H) = 47 \text{ MeV at } 1.4 \text{ TeV},$$

$$\Delta(m_H) = 44 \text{ MeV at } 3 \text{ TeV}.$$

A combination of both energy stages would lead to a precision of 32 MeV.



FCCEe Recoil Mass




Higgs boson width

Does the Higgs boson decay as predicted?

Γ_H at LHC



- Direct constraints from $m_{4\ell}$, $m_{\gamma\gamma}$ mass shape & $c\tau$ measurements: challenging even at HL-LHC
- Indirect constraint from interference between Higgs production and backgrounds:
 - ➔ between $gg \rightarrow H \rightarrow \gamma\gamma^*$ and $gg \rightarrow \gamma\gamma^*$ (statistics dominated)
 - ➔ high-mass $gg \rightarrow H \rightarrow VV$ and $gg \rightarrow VV$



At high-mass, off-shell Higgs production and non-resonant $gg \rightarrow VV$ background (box diagram) sizeable and negative interference

ATLAS projection for 3000 fb^{-1} at HL-LHC

$$\Gamma_H = 4.2^{+1.5}_{-2.1} \text{ MeV (stat+sys)}$$





Γ_H at Lepton Collider

[arXiv:1608.07538](https://arxiv.org/abs/1608.07538)

[arXiv:1710.07621](https://arxiv.org/abs/1710.07621)



Use $e^+e^- \rightarrow ZH$, reconstruct $Z \rightarrow \mu^+\mu^-$, e^+e^- :
reconstruct recoil mass (independent of Higgs final state)




CLIC350: $\Delta(\Gamma_H) = 6.7\%$

ILC250: $\Delta(\Gamma_H) = 3.9 - 2.5\%$ (depending on fit model)

ILC500: $\Delta(\Gamma_H) = 1.7 - 2.6\%$ (depending on fit model)

$e^+e^- \rightarrow ZH$, $Z \rightarrow q\bar{q}$, $H \rightarrow \text{invisible}$



CLIC350: $\Gamma_{\text{invis}}/\Gamma_H < 0.01$ at 90% C.L.

ILC500: $\Gamma_{\text{invis}}/\Gamma_H < 0.003$ at 95% C.L.

Rare Higgs Branching Ratios


Does the Higgs boson decay to second generation fermions?

Higgs Hadronic BRs at CLIC

arXiv:1608.07538



- Aim: resolve $H \rightarrow 2$ jets signal into $H \rightarrow b\bar{b}$, $H \rightarrow c\bar{c}$ and $H \rightarrow gg$
- Fit to multivariate-derived templates using flavour tagging info e.g. at 350 GeV, $HZ \rightarrow 4$ jets



With 500/fb at 350 GeV

Also with:

- ➔ $HZ \rightarrow 2$ jets + 2 ℓ ,
- ➔ $\nu\nu H \rightarrow E + 2$ jets

Decay	Statistical uncertainty	
	Higgsstrahlung	WW-fusion
$H \rightarrow b\bar{b}$	0.86 %	1.9 %
$H \rightarrow c\bar{c}$	14 %	26 %
$H \rightarrow gg$	6.1 %	10 %



$H \rightarrow c\bar{c}$ Search at LHCb

LHCb-CONF-2016-006



Search for VH , $V \rightarrow \ell$, $H \rightarrow b\bar{b}$, $c\bar{c}$ in forward production ($2 < \eta < 5$)

With 1.92 fb^{-1} , $H \rightarrow b\bar{b} < 50 \times \text{SM}$; $H \rightarrow c\bar{c} < 7900 \times \text{SM}$, (95% CL)



Very rough information projection: LHCb could reach $5 \times \text{SM}$ with 300 fb^{-1}

$H \rightarrow J/\psi \gamma$ at ATLAS




- Sensitive to Higgs coupling to charm quark
 - SM expectation: $\text{BR}(H \rightarrow J/\psi \gamma) = (2.9 \pm 0.2) \times 10^{-6}$
 - use $J/\psi \rightarrow \mu^+ \mu^-$ decay mode
 - $Z \rightarrow J/\psi \gamma$ as a cross check
- Using multivariate analysis, events in $m(\mu^+ \mu^- \gamma) \in 115\text{-}135 \text{ GeV}$: **~3** signal, **1700** background

Expected limits at 95% CL (using multivariate analysis):

- $\text{BR}(H \rightarrow J/\psi \gamma): (44^{+19}_{-12}) \times 10^{-6}$
- $\sigma(\text{gg} \rightarrow H) \times \text{BR}(H \rightarrow J/\psi \gamma): (3.1^{+0.9}_{-1.3}) \text{ fb}$

(no background systematics considered)



Also possible for lighter vectors:

Channel	Final state	Exp. SM $\times 10^{-6}$	Run2 limit
$H \rightarrow \rho \gamma$	$\pi^+ \pi^- \gamma$	16.8 ± 0.8	880
$H \rightarrow \phi \gamma$	$K^+ K^- \gamma$	2.31 ± 0.11	480

[arXiv:1712.02758](https://arxiv.org/abs/1712.02758)



$H \rightarrow \mu^+ \mu^-$ at HL-LHC



[CMS-TDR-17-001](#)

[ATLAS-TDR-026](#)

[PhysRevLett.119.051802](#)




- SM prediction is $\text{BR}(H \rightarrow \mu\mu) = 2.19 \times 10^{-4}$
- Run 1 limit is $2.8 \times \text{SM}$
- ATLAS production with 3000 fb⁻¹:
 - Observation at $\sim 7\sigma$
 - uncertainty of 20-25 % on signal strength

$H \rightarrow \mu^+ \mu^-$ at CLIC





$$\frac{\Delta[\sigma(H\nu_e\bar{\nu}_e) \times BR(H \rightarrow \mu^+\mu^-)]}{\sigma(H\nu_e\bar{\nu}_e) \times BR(H \rightarrow \mu^+\mu^-)} = 38\% \text{ at } 1.4 \text{ TeV},$$

$$\frac{\Delta[\sigma(H\nu_e\bar{\nu}_e) \times BR(H \rightarrow \mu^+\mu^-)]}{\sigma(H\nu_e\bar{\nu}_e) \times BR(H \rightarrow \mu^+\mu^-)} = 25\% \text{ at } 3 \text{ TeV}.$$



Very clean, narrow signal!

$H \rightarrow \mu^+ \mu^-$ & $H \rightarrow Z\gamma$ at FCCpp




Spin & CP

Only pseudoscalar in there?

Many places to look ... several different
parameterisations of CP used

CP Studies at HL-LHC

- $H \rightarrow ZZ \rightarrow 4\ell$ used to reconstruct the full angular decay structure.
- Sensitive to non-SM ($\mathbf{CP} = \mathbf{0}^+$) contributions.



$$A(H \rightarrow ZZ) = v^{-1} \left(\underbrace{a_1 m_Z^2 \epsilon_1^* \epsilon_2^*}_{\text{SM tree processes}} + \underbrace{a_2 f_{\mu\nu}^{*(1)} f^{*(2),\mu\nu}}_{\text{loop CP-even contributions}} + \underbrace{a_3 f_{\mu\nu}^{*(1)} \tilde{f}^{*(2),\mu\nu}}_{\text{CP-odd contributions (BSM)}} \right)$$

- Fit fraction of event (f_{ai}) and phases (ϕ_i) to observed decay:


$$\phi_{a_i} = \arg \left(\frac{a_i}{a_1} \right) \quad f_{a_i} = \frac{|a_i|^2 \sigma_i}{|a_1|^2 \sigma_1 + |a_i|^2 \sigma_i}$$

CP Studies at HL-LHC




ATL-PHYS-PUB-2013-013

CMS-PAS-FTR-16-002



$$a_1 = g_1 \frac{m_V^2}{m_H^2} + \frac{s}{m_H^2} \left(2g_2 + g_3 \frac{s}{\Lambda^2} \right); \quad a_2 = - \left(2g_2 + g_3 \frac{s}{\Lambda^2} \right); \quad a_3 = -2g_4,$$

$$f_{g_i} = \frac{|g_i|^2 \sigma_i}{|g_1|^2 \sigma_1 + |g_2|^2 \sigma_2 + |g_4|^2 \sigma_4}; \quad \phi_{g_i} = \arg \left(\frac{g_i}{g_1} \right).$$



Percent-level sensitivity, but still statistics limited

CP Studies at ILC250

$$\Delta\mathcal{L}_{hZZ} = \frac{1}{2} \frac{\tilde{b}}{v} h Z_{\mu\nu} \tilde{Z}^{\mu\nu}.$$

$\tilde{b} \equiv a_3$ can be measured to 0.5% at ILC250

$H \rightarrow \tau^+ \tau^-$ decay probed

$$\Delta\mathcal{L}_{h\tau\tau} = -\frac{\kappa_\tau y_\tau}{\sqrt{2}} h \tau^+ (\cos \phi + i \sin \phi \gamma_5) \tau^-$$

CP phase: $\phi=0$ in the SM
 ϕ measured to 3.8° at ILC250

CP Studies at FCCee



$$\Delta\mathcal{L}_{h\tau\tau} = -\frac{\kappa_\tau y_\tau}{\sqrt{2}} h\tau^+ (\cos\phi + i \sin\phi \gamma_5) \tau^-$$

➔ **$H \rightarrow \tau\tau$ decay is promising channel to study CP violation**


- Tree level couplings to quarks and leptons
- CP-even and CP-odd couplings induced at the same order

➔ **CP violation can be probed through τ polarization**

- τ decays clean enough that the spin information is not washed out by hadronization effects
- pion emission preferred in the direction of the τ spin in rest frame $\tau^\pm \rightarrow \rho^\pm \nu_\tau \rightarrow \pi^\pm \pi^0 \nu_\tau$
- exploring $\mathcal{L}_{hff} \propto h\bar{f}(\cos\Delta + i\gamma_5 \sin\Delta)f$
- model using effective lagrangian

Andres Rios (MIT), Aram Apyan (FNAL)

following arXiv:1308.1094





- 920 signal event in 5ab-1
- expected 68% CL
 - ❖ 0.17 radian (0.05 in GEN level study)
 - ❖ 9.7 degree (2.9 in GEN level study)

CP Studies at LHeC

U. Klein at HL/HE LHC meeting April 2018

LHeC CDR


LHeC provides access to CP structure of WWH and ttH coupling




$$\Gamma_{\mu\nu}^{(\text{BSM})}(p, q) = \frac{-g}{M_W} [\lambda (p \cdot q g_{\mu\nu} - p_\nu q_\mu) + i \lambda' \epsilon_{\mu\nu\rho\sigma} p^\rho q^\sigma]$$

CP-conserving
CP-violating

LHeC allows access to CP structure of WWH and ttH coupling




$$\mathcal{L} = -\frac{m_t}{v} \bar{t} [\kappa \cos \zeta_t + i\gamma_5 \sin \zeta_t] t h$$




LHeC


rapidity difference (H, \bar{t})



top polarisation



fiducial incl. cross-section



Differential Cross Sections




Is the devil in the detail?

Differential Cross Sections

Thomas Klinjsma, HL/HE
LHC Meeting, 8th April 2018



Transverse momentum $p_T(H)$

- ➔ Sensitivity to modifications of effective Higgs Yukawa couplings
- ➔ Sensitivity to finite top mass effects





Jet multiplicity N_{jets} & p_T of the first jet

- ➔ New physics in the loop, sensitivity at high p_T



Differential Cross Sections at HL-LHC





Personal Conclusions

- Lots still to learn about the Higgs boson: All possible future colliders will tell us much more about Higgs properties!
- If you (only) want measure Higgs properties & couplings soon, an e^+e^- collider is the best option.
- $\sqrt{s} = 250 \text{ GeV}$ (ILC250, FCCee): ZH production \Rightarrow good
- $\sqrt{s} = 380 \text{ GeV}$ (CLIC380, FCCee): adds $WW \rightarrow \nu\bar{\nu}H$ production \Rightarrow better
- $\sqrt{s} > 550 - 600 \text{ GeV}$ (ILC upgrade?, CLIC1500)
adds $t\bar{t}H$ and ZHH production \Rightarrow best

Backup

Higgs Couplings


Precise, model-independent measurements of couplings to fermions, W , Z .



e^+e^- collider at 250 - 380 GeV would be a Higgs Factory!
at higher collisions energies can test Higgs-self-coupling

CLIC Higgs Couplings Precision


[arXiv:1608.07538](https://arxiv.org/abs/1608.07538)



What kind of Higgs?

	Model	$b\bar{b}$	$c\bar{c}$	gg	WW	$\tau\tau$	ZZ	$\gamma\gamma$	$\mu\mu$
1	MSSM [38]	+4.8	-0.8	- 0.8	-0.2	+0.4	-0.5	+0.1	+0.3
2	Type II 2HD [39]	+10.1	-0.2	-0.2	0.0	+9.8	0.0	+0.1	+9.8
3	Type X 2HD [39]	-0.2	-0.2	-0.2	0.0	+7.8	0.0	0.0	+7.8
4	Type Y 2HD [39]	+10.1	-0.2	-0.2	0.0	-0.2	0.0	0.1	-0.2
5	Composite Higgs [40]	-6.4	-6.4	-6.4	-2.1	-6.4	-2.1	-2.1	-6.4
6	Little Higgs w. T-parity [41]	0.0	0.0	-6.1	-2.5	0.0	-2.5	-1.5	0.0
7	Little Higgs w. T-parity [42]	-7.8	-4.6	-3.5	-1.5	-7.8	-1.5	-1.0	-7.8
8	Higgs-Radion [43]	-1.5	- 1.5	+10.	-1.5	-1.5	-1.5	-1.0	-1.5
9	Higgs Singlet [44]	-3.5	-3.5	-3.5	-3.5	-3.5	-3.5	-3.5	-3.5

Table 3: Percent deviations from SM for Higgs boson couplings to SM states in various new physics models. These model points are unlikely to be discoverable at 14 TeV LHC through new particle searches even after the high luminosity era (3 ab^{-1} of integrated luminosity). From [20].



Indirect constraint on Γ_H from offshell production

- $\sigma_{\text{offshell}} \sim g_g^2 g_V^2$ does not depend on the total width Γ_H , σ_{onshell} does
 - In terms of coupling modifiers

$$\frac{\sigma_{\text{off-shell}}^{gg \rightarrow H^* \rightarrow ZZ}}{\sigma_{\text{off-shell, SM}}^{gg \rightarrow H^* \rightarrow ZZ}} = \mu_{\text{off-shell}} = \kappa_{g,\text{off-shell}}^2 \cdot \kappa_{V,\text{off-shell}}^2 \quad \frac{\sigma_{\text{on-shell}}^{gg \rightarrow H \rightarrow ZZ}}{\sigma_{\text{on-shell, SM}}^{gg \rightarrow H \rightarrow ZZ}} = \mu_{\text{on-shell}} = \frac{\kappa_{g,\text{on-shell}}^2 \cdot \kappa_{V,\text{on-shell}}^2}{\Gamma_H / \Gamma_H^{\text{SM}}}$$

- Under the assumption of equal on-peak and off-peak coupling modifiers, limit on μ_{offshell} can be **reinterpreted**, combined **with** μ_{onshell} , as **limit on Γ_H**
 - Strong assumption, $\kappa_g(s)$ sensitive to possible new physics at higher mass scales
 - New physics which modify off-shell signal strength do not change bkg predictions

$$\kappa_{g,\text{on-shell}}^2 \kappa_{V,\text{on-shell}}^2 \leq \kappa_{g,\text{off-shell}}^2 \kappa_{V,\text{off-shell}}^2$$

- Latest experimental results (WW+ZZ in Run1 for ATLAS and CMS, 4l Run2 CMS):

$\Gamma_H < 22.7 \text{ MeV @ 95\%CL}$
($<33 \text{ MeV exp.}$)

ATLAS Run1 Eur.Phys.J.C (2015) 75:335

$\Gamma_H < 13 \text{ MeV @ 95\%CL}$
($<26 \text{ MeV exp.}$)

CMS Run1 JHEP 09 (2016) 051

4l: $\Gamma_H < 41 \text{ MeV @ 95\%CL}$
($<32 \text{ MeV exp.}$)

CMS Run2, 12.9 fb⁻¹ CMS PAS HIG-16-033

- For HL-LHC most of the consideration done for μ_{offshell} valid here as well
 - In this interpretation, the uncertainty on $\mu_{\text{off-shell}}$ dominates
 - $\sim 5\%$ precision achievable for $\mu_{\text{onshell}}^{ZZ}$
 - Estimate using 4l alone by ATLAS (10% syst on $R_{H^*}^B$)

$\Gamma_H = 4.2^{+1.5}_{-2.1} \text{ MeV}$


ATL-PHYS-PUB-2015-024



LHC: $H \rightarrow c\bar{c}$



- * Search for leptonic W/Z+ $H \rightarrow b\bar{b}/c\bar{c}$ final states (Forward production)
- * LHCb angular acceptance \rightarrow Challenging for integrated cross sections and uncertainties



300 fb⁻¹ at 14 TeV:

$$|\kappa_c| \lesssim 7$$

30% di-c-tagging efficiency:

$$|\kappa_c| \lesssim 4$$

better electron reconstruction:

$$|\kappa_c| \lesssim 3$$

further improvements:

$$|\kappa_c| \lesssim 2.2$$

projections taken from talk by Mike Williams

- * First publication $H \rightarrow c\bar{c}$ from LHCb \rightarrow Possible reach 5 xSM @ 300 fb⁻¹
- * Extrapolation $H \rightarrow J/\Psi \gamma$ @ HL-LHC : 15 xSM @ 3000 fb⁻¹ (ATLAS)
- * $ZH \rightarrow c\bar{c}$ @ Run-2 (arxiv:1802.04329): $\mu < 110 (150^{+80}_{-40})$ @ 95 % C.L. (ATLAS)