

**Improved analysis on  $\gamma\gamma \rightarrow \text{higgs} \rightarrow b\bar{b}$**   
**for SM and MSSM,**  
**with overlaying events,**  
**vertex smearing and crab crossing**

P. Niezurawski, A. F. Żarnecki, M. Krawczyk

Faculty of Physics  
Warsaw University

# Overview

Analysis of  $\sigma(\gamma\gamma \rightarrow higgs \rightarrow b\bar{b})$  measurement  
Montpellier:

- realistic  $\gamma\gamma$ -spectra
- $b$ -tagging
- overlaying events  $\gamma\gamma \rightarrow hadrons$  (OE)

⇒ results for SM at  $M_h = 120, 130, 140, 150, 160$  GeV

⇒ results for MSSM at  $M_A = 200, 250, 300, 350$  GeV  
with  $\tan\beta = 7, M_2 = \mu = 200$  GeV (following M. Mühlleitner *et al.*)

**NEW:**

- improved treatment of overlaying events
- crossing angle
- primary vertex distribution
- $\gamma\gamma \rightarrow W^+W^-$  background contribution

$$\gamma\gamma \longrightarrow \text{higgs} \longrightarrow b\bar{b}$$

Photon-photon spectrum: CompAZ

Signal: HDECAY, PYTHIA

Background: NLO  $Q\bar{Q}(g)$  (G. Jikia);  $\gamma\gamma \rightarrow W^+W^-$  (PYTHIA)

Pile-up events  $\gamma\gamma \rightarrow \text{hadrons}$  (PYTHIA) with realistic  $\gamma\gamma$ -luminosity spectrum (V. Telnov)

Parton Shower (signal &  $\gamma\gamma \rightarrow W^+W^-$ ): PYTHIA

Fragmentation: PYTHIA (Lund)

Detector performance: SIMDET 4.01

Jets: Durham algorithm,  $y_{cut} = 0.02$  (clusters & tracks with  $|\cos\theta| > \cos\theta_{det} = 0.85$  ignored)


**Selection of  $b\bar{b}$  events for  $M_{higgs} = 120$  (300) GeV:**

- OE-jets suppression: consider only jets with  $p_T^{jet}/E_T > 0.1$
- $N_{jets} = 2, 3$  (2)
- $|\cos\theta_{jet}| < 0.71$  (0.65) for each jet
- $|P_z|/E < 0.12$  (0.07) where  $P_z = \sum p_z^{jet}$  and  $E = \sum E^{jet}$
- To suppress  $\gamma\gamma \rightarrow W^+W^-$  background, for  $M_{higgs} > 160$  GeV (MSSM):  
 $P_T/E_T < 0.11$  &  $M_{jet1} < 70$  GeV &  $M_{jet2} < 50$  GeV; at least 7 tracks in each jet
- ZVTOP-B-Hadron-Tagger by T. Kuhl
- Correction for crossing angle: boost with  $\beta = \sin(\alpha_c/2)$




# $\gamma\gamma \rightarrow \text{hadrons events}$

Angular  $E_T$ -flow per bunch crossing.



Generation for  $\sqrt{s_{ee}} = 210.5$  GeV.

Angular cuts



Detector mask

Particles on Pythia level:  $\cos \theta_{\text{mask}} = 0.99$

OE suppression

Tracks & clusters:  $\cos \theta_{\text{det}} = 0.85$

$\gamma\gamma \rightarrow Q\bar{Q}(g)$  suppression



Jets:  $\cos \theta_{\text{jet}} = 0.71$  (0.65)

# Transverse momenta of jets

Discriminating between gluon-jets (PS) & OE

SM - signal only

MSSM - signal only



Contribution of OE  $\Rightarrow$  additional cut:  $p_T^{\text{jet}} / E_T > 0.1$  for each jet

# Crab-wise crossing of beams


$$\sigma'_x = \sqrt{\frac{1}{2}(\sigma_x^2 + \sigma_z^2 \tan^2(\alpha_c/2))}$$

$$\sigma'_y = \sigma_y/\sqrt{2}$$

$$\sigma'_z = \sigma_z/\sqrt{2}$$

Bunch:  $\sigma_x = 140 \text{ nm}$   $\sigma_y = 15 \text{ nm}$   $\sigma_z = 0.3 \text{ mm}$

Primary vertex:  $\sigma'_x = 3.6 \text{ }\mu\text{m}$   $\sigma'_y = 11 \text{ nm}$   $\sigma'_z = 0.2 \text{ mm}$



$$\alpha_c = 34 \text{ mrad}$$



# *higgs-tagging* at $M_h = 120 \text{ GeV}$

*higgs-tagging*: a cut on the ratio  
of  $\gamma\gamma \rightarrow h \rightarrow b\bar{b}$

to  $\gamma\gamma \rightarrow b\bar{b}(g), c\bar{c}(g)$  events

$\Rightarrow \varepsilon_{higgs} = 70 \%$

$\varepsilon_{bb} = 66\%, \varepsilon_{cc} = 4\%$

Earlier we used *b*-tagging:

a cut on the ratio


of  $\gamma\gamma \rightarrow b\bar{b}(g)$

to  $\gamma\gamma \rightarrow c\bar{c}(g)$  events

$\Rightarrow \varepsilon_{higgs} = 85 \%$

$\varepsilon_{bb} = 82\%, \varepsilon_{cc} = 2\%$

Tighter cuts are needed  
due to OE contribution




Black stars – optimized selection

Black+white stars – analysis without OE


# SM, $M_h = 120$ GeV

Corrected reconstructed mass:  $W_{\text{corr}} \equiv \sqrt{W_{\text{rec}}^2 + 2P_T(E + P_T)}$   
 Acta Phys. Pol. B34 177-187 2003, hep-ph/0208234

OLD



NEW





1.6%  $\rightarrow$  2%



# SM, $M_h = 120-160$ GeV

NEW:  $\gamma\gamma \rightarrow W^+W^-$  background

Precision of  $\gamma\gamma \rightarrow h \rightarrow b\bar{b}$  measurement




$M_h = 160$  GeV



# MSSM, $M_A = 300$ GeV

NEW


## Optimal *higgs*-tagging




$$\frac{\#\gamma\gamma \rightarrow h \rightarrow b\bar{b}}{\#\gamma\gamma \rightarrow b\bar{b}(g), c\bar{c}(g)}$$

$$\varepsilon_{higgs} = 68\%$$

Number of overlaying events:  $\sim 2$  per bc



# MSSM, $M_A = 200-350$ GeV



Precision between 11% and 23%.

# Conclusions

## NEW analyses for SM and MSSM:

- High precision for measurement of SM & MSSM higgses despite  $\gamma\gamma \rightarrow \text{hadrons}$  pile-up events and primary vertex distribution
- Cut on  $p_T^{\text{jet}} / E_T$  improves rejection of OE jets
- *higgs-tagging*: cut on the ratio of  $\gamma\gamma \rightarrow h \rightarrow b\bar{b}$  to  $\gamma\gamma \rightarrow b\bar{b}(g), c\bar{c}(g)$  events  $\Rightarrow$  improved  $\gamma\gamma \rightarrow Q\bar{Q}(g)$  background suppression
- Precision of 2% for  $\Gamma(h \rightarrow \gamma\gamma)\text{Br}(h \rightarrow b\bar{b})$  at  $M_h = 120$  GeV
- Precision between 11% and 23% for  $M_A = 200\text{--}350$  GeV.
- Inclusion of  $\gamma\gamma \rightarrow W^+W^-$  background influences precision estimates for MSSM case. Improvement in discriminating of  $W^+W^-$  possible

### Plans:

- MSSM: parameters space scan

