




# Particle Physics & the Energy Frontier Beyond HL-LHC

*Ian Shipsey*  
*Oxford*

*PPAP 2018*  
*July 16-17 2018*  
*RAL*



# OUTLINE

What are the important questions to consider at the Energy Frontier in preparing the UK input to the European Strategy Update?

Particle Physics circa 2018

Opportunities for *great discoveries*  
*@ the energy frontier*



# Will present information that will be relevant to considering this recommendation


## European Strategy Workshop, IPPP 16-18th April 2018: Meeting Summary

Overview (J. Evans, S. Farrington, E. Goudzovski, M. Patel, M. Spannowsky)

4) The physics cases for FCC and CLIC are clearly both strong but there are resource implications in pushing both R&D programs forward during the 2020s. The last UK ES submission said that *“a timely decision should be taken on optimal next-generation collider facilities for exploitation of LHC discoveries”*. The final 2013 ES update document said *“to stay at the forefront of particle physics, Europe needs to be in a position to propose an ambitious post-LHC accelerator project at CERN by the time of the next Strategy update, when physics results from the LHC running at 14 TeV will be available”*. It is the recommendation of the organisers of this workshop that it should be considered, in a UK community meeting, whether a decision can now be made on a definitive UK recommendation. If a consensus cannot be reached, then it could be debated in the community meeting whether to put forward to the ES process that its committee makes a definitive recommendation by 2020.

2012.7.4


# discovery of Higgs boson



theory : 1964

design : 1984

construction : 1998



The Higgs enables  
atoms to exist

# Outstanding Questions in Particle Physics *circa 2011*

## EWSB

- ☐ Does the Higgs boson exist?

## Quarks and leptons:

- ☐ why 3 families ?
- ☐ masses and mixing
- ☐  $CP$  violation in the lepton sector
- ☐ matter and antimatter asymmetry
- ☐ baryon and charged lepton number violation

## Physics at the highest E-scales:

- ☐ how is gravity connected with the other forces ?
- ☐ do forces unify at high energy ?

## Dark matter:

- ☐ composition: WIMP, sterile neutrinos, axions, other hidden sector particles, ..
- ☐ one type or more ?
- ☐ only gravitational or other interactions ?

## The two epochs of Universe's accelerated expansion:

- ☐ primordial: is inflation correct ?  
which (scalar) fields? role of quantum gravity?
- ☐ today: dark energy (why is  $\Lambda$  so small?) or gravity modification ?

## Neutrinos:

- ☐  $\nu$  masses and their origin
- ☐ what is the role of  $H(125)$  ?
- ☐ Majorana or Dirac ?
- ☐  $CP$  violation
- ☐ additional species  $\rightarrow$  sterile  $\nu$  ?

# Outstanding Questions in Particle Physics *circa 2018*

... there has never been a better time to be a particle physicist!

## Higgs boson and EWSB

- ☐  $m_H$  natural or fine-tuned ?  
→ if natural: what new physics/symmetry?
- ☐ does it regularize the divergent  $V_L V_L$  cross-section at high  $M(V_L V_L)$  ? Or is there a new dynamics ?
- ☐ elementary or composite Higgs ?
- ☐ is it alone or are there other Higgs bosons ?
- ☐ origin of couplings to fermions
- ☐ coupling to dark matter ?
- ☐ does it violate CP ?
- ☐ cosmological EW phase transition

## Dark matter:

- ☐ composition: WIMP, sterile neutrinos, axions, other hidden sector particles, ..
- ☐ one type or more ?
- ☐ only gravitational or other interactions ?

## The two epochs of Universe's accelerated expansion:

- ☐ primordial: is inflation correct ?  
which (scalar) fields? role of quantum gravity?
- ☐ today: dark energy (why is  $\Lambda$  so small?) or gravity modification ?

## Quarks and leptons:

- ☐ why 3 families ?
- ☐ masses and mixing
- ☐ CP violation in the lepton sector
- ☐ matter and antimatter asymmetry
- ☐ baryon and charged lepton number violation

## Physics at the highest E-scales:

- ☐ how is gravity connected with the other forces ?
- ☐ do forces unify at high energy ?

## Neutrinos:

- ☐  $\nu$  masses and their origin
- ☐ what is the role of  $H(125)$  ?
- ☐ Majorana or Dirac ?
- ☐ CP violation
- ☐ additional species → sterile  $\nu$  ?



These questions are compelling, difficult and intertwined → require multiple approaches  
 high-E colliders, neutrino experiments (solar, short/long baseline, reactors  
 $0\nu\beta\beta$  decays), cosmic surveys (CMB, optical/IR spectroscopic and photometric ), dark matter  
 direct, indirect and astrophysical detection, precision measurements of quark and lepton  
 rare decays and phenomena, dedicated searches (axions, dark-sector particles), ...

### Main questions and main approaches to address them

	High-E colliders	High-precision experiments	Neutrino experiments	Dedicated searches	Cosmic surveys
Higgs , EWSB	x				
Neutrinos			x	x	x
Dark Matter	x			x	x
Flavour, CP-violation	x	x	x	x	
New particles and forces	x	x	x	x	
Universe acceleration					x

These complementary approaches are ALL needed: their combination is crucial to explore the largest range of E scales, properly interpret signs of new physics, and build a coherent picture of the underlying theory.



# Standard Model Lagrangian

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + i \bar{\psi} \not{D} \psi + h$$

Describes everything experimentally confirmed before 2012

Higgs sector

$$+ \sum_i \bar{\psi}_i \gamma_{ij} \psi_j \phi + h$$

$$+ |D_\mu \phi|^2 - V(\phi)$$

Yukawa coupling with new scalar (completely new interaction type)  
ttH,  $H \rightarrow b\bar{b}$  and  $H \rightarrow \tau\bar{\tau}$  are important !

Higgs potential ( $\mu^2 \phi^2 + \lambda \phi^4$ )


Gauge boson interaction with new scalar (new for scalar, but known for fermions)




PPAP 16/7/18 -- I. Shipsey



Our work has the potential to lead to a reconciliation of the two great edifices of physics



General Relativity



Quantum Mechanics

# No-lose completion of the Standard Model

Guaranteed  
discoveries

W & Z	CERN SppS (1983)
Top quark	Tevatron (1995)
Higgs	LHC (2012)

# No-lose completion of the Standard Model

Now that the Standard Model is complete,  
there are no further no-lose theorems  
In principle, the Standard Model could be  
valid to the Planck scale

No guaranteed  
discoveries



There are no guaranteed discoveries

Higgs is central to SM & BSM  
& a guaranteed deliverable  
@ any future collider

Direct  
High E  
pp (later)

Precision  
Lower E  
ee (sooner)


Focus on Higgs @ ee  
& in particular the couplings

# IMPACT of 125 GeV on Energy Frontier

The low mass of the Higgs makes  $e^+e^-$  Higgs factories both linear and circular tractable & has consequently modified & simplified the landscape of accelerator options at the energy frontier since 2012

Higgs	ILC 250	Energy	ILC
Factories	CLIC 380	Frontier	CLIC
(also Z,	CEPC		SPPC
W, t)	FCC-ee		FCC-pp
			FCC-eh
	LHeC		


# Higgs @ a pp collider



LHC –

- large Higgs cross section
- 150M Higgs per exp.
- dominated by QCD events
- $S(H)/B(\text{All}) \sim 10^{-10}$
- Pile-up and jet overlap
- Not knowing

$(P_H, E_H)_{\text{initial}}$





Precisions on H couplings

~(5-10)%  
expected





Key feature:

Higgs coupling depends on the particle mass



# $e^+e^-$ Higgs production @ 250 GeV



$$m_{\text{recoil}}^2 = (\sqrt{s} - E_{f\bar{f}})^2 - p_{f\bar{f}}^2 = s - 2E_{f\bar{f}}\sqrt{s} + m_{f\bar{f}}^2$$

The tagging of  $e^+e^- \rightarrow ZH$  events through the recoil mass method is independent of the Higgs boson decay.

# $e^+e^-$ Higgs production @ 250 GeV

Higgs events are readily isolated from background.  
All standard Higgs decay modes are visible.

Measurement accuracies are such that 1% coupling measurements are feasible.

The absolute cross section for  $e^+e^- \rightarrow Zh$  can be measured.


At 250 GeV, to first approximation, any Z boson with  $E_{lab} = 110$  GeV is recoiling against a Higgs boson.



# ILC/CLIC and CEPC/FCC(ee)


Linear accelerator can reach high energies ~multi-TeV  
with high luminosity

- Can avoid synchrotron radiation
- High accelerating field to achieve high energy
  - ✓ Normal conducting accelerating structures (CLIC)
- High beam current and quality to achieve the luminosity
  - ✓ High quality of components
  - ✓ Nano beams




Circular accelerator can reach high luminosity  
at lower energies

- Can store and re-collide the beams
- Experience
- Synchrotron radiation limits the energy and beam quality




# future lepton collider luminosities



→ earliest possible physics starting dates

- ILC250: 2032
- CLIC350: 2035
- FCC-ee: 2039
- CEPC: 2030

# Higgs mass



$$\delta m_h \sim 300 \text{ MeV}$$


➤ Higgs mass precision can be limitation of coupling fit precision

$$\delta_W = 6.9 \cdot \delta m_h, \quad \delta_Z = 7.7 \cdot \delta m_h$$

[Almeida, Lee, Pokorski, Wells '13]

➤ through leptonic recoil in  $Z \rightarrow \mu^+ \mu^-$   
the Higgs mass can be constrained  
to 14 MeV [LCC Physics Working Group '18]

➤ impact on Z/W couplings  $\sim 0.1\%$



# Higgs Branching Fraction Measurements

With both  $\sigma_{Zh}$  and  $\sigma_{Zh} \cdot \text{BR}$  measured, the absolute branching ratios can be determined independently

Decay mode	$\sigma(ZH) \times \text{BR}$	BR
$H \rightarrow b\bar{b}$	0.28%	0.57%
$H \rightarrow c\bar{c}$	2.2%	2.3%
$H \rightarrow gg$	1.6%	1.7%
$H \rightarrow \tau\tau$	1.2%	1.3%
$H \rightarrow WW$	1.5%	1.6%
$H \rightarrow ZZ$	4.3%	4.3%
$H \rightarrow \gamma\gamma$	9.0%	9.0%
$H \rightarrow \mu\mu$	17%	17%
$H \rightarrow \text{inv}$	—	0.28%

Relative error (%) CEPC Pre-CDR

Most precise:  $\text{BR}_{bb}$  and  $\text{BR}_{\tau\tau}$ , ILC (CEPC) 0.89% (0.57%) and 1.4% (1.3%) respectively. If there are O(1%) or larger exotic decay modes, a first hint would be provided by observing the resulting deviations in  $\text{BR}_{bb}$  and  $\text{BR}_{\tau\tau}$ .

# Measuring the Higgs width

$$BR(h \rightarrow A\bar{A}) = \Gamma(h \rightarrow A\bar{A})/\Gamma_h$$

$\Gamma(H_{125})_{SM} = 4.1 \text{ MeV}$ , too small to be measured directly determine indirectly;  
requires a formalism.

Traditionally width is determined using the  $\kappa$  parametrization.

Assumes Higgs coupling to A is modified from SM value by a multiplicative factor  $\kappa_A$

$$\frac{\Gamma(h \rightarrow ZZ^*)}{SM} = \kappa_Z^2, \quad \frac{\sigma(e^+e^- \rightarrow Zh)}{SM} = \kappa_Z^2$$

$$\Gamma_H = \frac{\Gamma(H \rightarrow ZZ^*)}{BR(H \rightarrow ZZ^*)} \propto \frac{\sigma(ZH)}{BR(H \rightarrow ZZ^*)}.$$


(CEPC)

$\Gamma_H$	$\sigma(ZH)$
3.3%	0.50%

( LHC limits  $\sim \times 3$  SM ATLAS  
14.4 MeV new ICHEP 18)



# Higgs Coupling Measurements comparison ILC & CEPC



$$\frac{\Gamma(h \rightarrow ZZ^*)}{SM} = \kappa_Z^2$$

ILC  
CEPC  
~1% uncertainty

$K_Z \sim 0.2\%$

$\delta \Gamma \sim 3\%$  (12 KeV) c.f. Current LHC limit  $\sim x3 \Gamma(SM)$


ILC and CepC achieve similar precision

# Higgs Coupling Measurements @ ILC & HL-LHC

## Effective Field Theory

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \sum_i \frac{c_i}{\Lambda^2} \mathcal{O}_i$$




	ILC250 2 ab <sup>-1</sup> w. pol.	ILC250+500 full ILC 250+500 GeV
$g(hb\bar{b})$	1.1	0.58
$g(hc\bar{c})$	1.9	1.2
$g(hgg)$	1.7	0.95
$g(hWW)$	0.67	0.34
$g(h\tau\tau)$	1.2	0.74
$g(hZZ)$	0.68	0.35
$g(h\gamma\gamma)$	1.2	1.0
$g(h\mu\mu)$	5.6	5.1
$g(hb\bar{b})/g(hWW)$	0.88	0.46
$g(hWW)/g(hZZ)$	0.07	0.05
$\Gamma_h$	2.5	1.6
$BR(h \rightarrow inv)$	0.32	0.29
$BR(h \rightarrow other)$	1.6	1.2



Physics Case for the 250 GeV Stage of the ILC, arXiv:1710.07621

(Standard Model Effective Field Theory (EFT) formalism.)

# Higgs couplings sensitivity to new physics



Full exploitation of a precision electron collider is the path to a model-independent measurement of the width and model-independent percent (or better) measurement of the couplings

ILC shown

Similar considerations apply to a circular e<sup>+</sup>e<sup>-</sup> collider


# Many BSM models impact Higgs couplings at percent level

Model	$b\bar{b}$	$c\bar{c}$	$gg$	$WW$	$\tau\tau$	$ZZ$	$\gamma\gamma$	$\mu\mu$
1 MSSM [38]	+4.8	-0.8	- 0.8	-0.2	+0.4	-0.5	+0.1	+0.3
2 Type II 2HD [39]	+10.1	-0.2	-0.2	0.0	+9.8	0.0	+0.1	+9.8
3 Type X 2HD [39]	-0.2	-0.2	-0.2	0.0	+7.8	0.0	0.0	+7.8
4 Type Y 2HD [39]	+10.1	-0.2	-0.2	0.0	-0.2	0.0	0.1	-0.2
5 Composite Higgs [40]	-6.4	-6.4	-6.4	-2.1	-6.4	-2.1	-2.1	-6.4
6 Little Higgs w. T-parity [41]	0.0	0.0	-6.1	-2.5	0.0	-2.5	-1.5	0.0
7 Little Higgs w. T-parity [42]	-7.8	-4.6	-3.5	-1.5	-7.8	-1.5	-1.0	-7.8
8 Higgs-Radion [43]	-1.5	- 1.5	+10.	-1.5	-1.5	-1.5	-1.0	-1.5
9 Higgs Singlet [44]	-3.5	-3.5	-3.5	-3.5	-3.5	-3.5	-3.5	-3.5

For the models shown above LHC not likely sensitive with full HL-LHC dataset

ILC250 Physics Case arXiv 1710.07621



# Many BSM models impact Higgs couplings at percent level



For the models shown above LHC not likely sensitive with full HL-LHC dataset

ILC250 Physics Case arXiv 1710.07621

# BSM physics through exotic Higgs decays



Uses ILC250 Physics Case arXiv 1710.07621

Z. Liu, H. Zhang, LT Wang, 1612.09284

PPAP 16/7/18 -- I. Shipsey




# Electroweak programme (W, Z) FCC-ee & CEPC (revisiting LEP with 100,000 times the Luminosity)

FCC-ee @ Z-pole  $3 \times 10^{12}$  Z bosons in 2 years

CEPC @ Z-pole  $10^{12}$  Z bosons in 2 years

Observable	LEP precision	CEPC precision	CEPC runs
$m_Z$	2 MeV	0.5 MeV	Z threshold scan
$A_{FB}^b$	1.7%	0.1%	Z threshold scan
$\sin^2 \theta_W^{\text{eff}}$	0.07%	0.002%	Z threshold scan
$R_b$	0.3%	0.02%	Z pole
$R_\mu$	0.2%	0.01%	Z pole
$N_\nu$	1.7%	0.05%	ZH runs
$m_W$	33 MeV	2-3 MeV	ZH runs
$m_W$	33 MeV	1 MeV	WW threshold

# Electroweak programme (W, Z) FCC-ee & CEPC




Current values  $\delta M_W$  :16 MeV Tevatron (comb.) 19 MeV ATLAS, SM Fit 4 MeV

# Future Circular Collider (FCC) – proton collider

## Higgs production

Compared to LHC at 14 TeV the cross section increases with a factor of about 16 at NNNLO. Together with a larger luminosity, one can expect 60-400x more events.



## Top Yukawa coupling

Measurement to 1% precision

## Higgs self-coupling

Measurement to 3-5% precision

## Higgs invisible decay Branching Ratio

Sensitivity down to  $3\text{-}5 \times 10^{-4}$

## Top quark production

Cross section increases x35 compared to LHC at 14 TeV, and might collect up to  $10^{12}$  top quarks


## New physics phenomena

In general direct sensitivity to processes with mass scales up to 10-40 TeV.


# Compact Linear Collider (CLIC)

CERN-2016-004

[arXiv:1608.07537](https://arxiv.org/abs/1608.07537)



CLIC aims at an acceleration gradient of 100 MV/m. A drive beam is decelerated in dedicated Power Extraction and Transfer Structures (PETS), and the generated RF power is transferred to the main beam.



# CLIC – some physics highlights

## Higgs & top quark characterization


Precision on top quark Yukawa of  $\sim 4\%$ ,  $m(\text{top})$  to 100 MeV (x5 better than HL-LHC) and Higgs self-coupling of  $\sim 20\%$ .

## Staged approach



First period as a Higgs/top factory, including a run at top quark pair threshold, thereafter operate at higher energies (upto 3TeV) which give access to:

- $t\bar{t}h$  and  $HHH$  couplings, as well as
- study any accessible new particles discovered @LHC
- searches for new phenomena


P. Burrows @ ICFA Seminar 2017



500 fb<sup>-1</sup> 380 GeV, 1.5 ab<sup>-1</sup> 1.5 TeV  
and 3 ab<sup>-1</sup> at 3 TeV.



PPAP 16/7/18 -- I. Shipsey



Total: 3 x 10<sup>6</sup> Higgs



## CLIC roadmap

### 2013 - 2019 Development Phase

Development of a Project Plan for a staged CLIC implementation in line with LHC results; technical developments with industry, performance studies for accelerator parts and systems, detector technology demonstrators

### 2020 - 2025 Preparation Phase

Finalisation of implementation parameters, preparation for industrial procurement, Drive Beam Facility and other system verifications, Technical Proposal of the experiment, site authorisation

### 2026 - 2034 Construction Phase

Construction of the first CLIC accelerator stage compatible with implementation of further stages; construction of the experiment; hardware commissioning

### 2019 - 2020 Decisions

Update of the European Strategy for Particle Physics; decision towards a next CERN project at the energy frontier (e.g. CLIC, FCC)


### 2025 Construction Start

Ready for construction; start of excavations

### 2035 First Beams

Getting ready for data taking by the time the LHC programme reaches completion

CLIC working on an implementation plan & cost reduction as input to European Particle Physics Strategy




# The 2013 European Particle Physics Strategy

*“There is a strong scientific case for an **electron-positron collider**, ... Europe looks forward to a proposal from Japan to discuss a possible participation.”*


Waiting for a statement from the Japanese Government for their willingness to host ILC before end of 2018

# International Linear Collider (ILC)



# Technology connection with the European XFEL at DESY

The 3.4 km long European XFEL generates extremely intense X-ray flashes to be used by researchers from all over the world.



First mass production in industry of SC radio frequency TESLA technology (from about 100 accelerator modules at the XFEL to about 2000 at ILC).

XFEL : 80% of the cavities reach a gradient of 33 MV/m

ILC : 90% of the cavities need a gradient of 35 MV/m

} Denis Kostin @ LCWS2017, Oct 24


This demonstrates the goal for the ILC is within reach.



# International Linear Collider (ILC) – 500 GeV → 250 GeV

Cost reduction both by scaling from 500 GeV to 250 GeV with a focus on Higgs physics, and by technological innovations on the superconducting materials (Nb) and cavity construction (surface process).

Physics Case for the 250 GeV Stage of the ILC, arXiv:1710.07621




ILC would be on a site surveyed for capability to reach 1 TeV – once ILC250 constructed extensions to at least 375 GeV would be almost guaranteed since we know that tt physics is very interesting and the enormous infrastructure/people investment for ILC250 would hardly be written off by Japan after 10 years.

PPAP 16/7/18 -- I. Shipsey





# Linear Collider detector & physics studies: Europe engaged



The LCC physics & detector directorate is responsible for activities that advance the physics and detectors of the linear collider.

## Three detector concepts:

- ILD: 71 institutions mostly from the European Region
- SiD: 24 institutions many from the European Region
- CLICdp: 29 institutions mostly from the European Region



## Three detector R&D groups:

- CALICE: 57 institutions mostly from the European Region
- LCTPC: 32 institutions many from the European Region
- ECAL: 14 institutions mostly from the European Region

# UK groups' interests in ILC/CLIC

All UK PP groups  
are represented  
in LCUK

~75 faculty have  
expressed interest  
in physics /  
detector /  
accelerator

UK expertise puts us  
in a strong position to  
play leading roles


*P. Burrows (LCUK)*

Technical system Institute	Accelerator					Detector			Physics
	BDS/MDI	DR	Beam dumps	e+ source	RF	Si tracker	Calorimetry	DAQ	
Birmingham	X					X	X		X
Bristol						X		X	X
Cambridge							X		X
STFC – Daresbury Laboratory	X			X	X				
Durham IPPP									X
Edinburgh						X			X
Glasgow						X			X
Imperial College							X	X	X
Lancaster				X	X	X			X
Liverpool		X				X			X
Manchester	X				X	X			X
Open University						X			
Oxford	X	X			X	X		X	X
QMUL						X			X
STFC – RAL			X			X			X
RHUL	X							X	X
Sheffield						X			X
Southampton									X
Sussex								X	X
UCL	X						X	X	X
Warwick						X			X

Many of these capabilities would also be of relevance to a future circular  
electron-positron collider should plans for either CEPC or FCCee proceed



# Future Circular Collider (FCC) Study




International FCC collaboration (CERN as host lab) to study:

- ***pp*-collider (*FCC-hh*)**  
→ main emphasis, defining infrastructure requirements

**~16 T  $\Rightarrow$  100 TeV *pp* 100 km**

- **~100 km tunnel infrastructure** in Geneva area, site specific
- ***e<sup>+</sup>e<sup>-</sup>* collider (*FCC-ee*)**, as potential first step
- **HE-LHC** with *FCC-hh* technology
- ***p-e* (*FCC-he*) option**, IP integration, *e<sup>-</sup>* from ERL



Physics Cases



Experiments




Collider Designs



R&D Programs



Infrastructures



Cost Estimates



# FCC : physics and performance targets

## FCC-ee:

- Exploration of 10 to 100 TeV energy scale via couplings with precision measurements
- ~20-50 fold improved precision on many EW quantities (equiv. to factor 5-7 in mass) ( $m_Z$ ,  $m_W$ ,  $m_{\text{top}}$ ,  $\sin^2 \theta_W^{\text{eff}}$ ,  $R_b$ ,  $\alpha_{\text{QED}}(m_Z)$ ,  $\alpha_s(m_Z)$ ,  $m_W m_\tau$ ), Higgs and top quark couplings)
- Machine design for highest possible luminosities at Z, WW, ZH and  $t\bar{t}$  working points

## FCC-hh:

- Highest center of mass energy for direct production up to 20 - 30 TeV
- Huge production rates for single and multiple production of SM bosons (H,W,Z) and quarks
- Machine design for 100 TeV c.m. energy & integrated luminosity  $\sim 20\text{ab}^{-1}$  within 25 years

## HE-LHC:

- Doubling LHC collision energy with FCC-hh 16 T magnet technology
- c.m. energy  $\sim 27\text{ TeV} = 14\text{ TeV} \times 16\text{ T}/8.33\text{ T}$ , target luminosity  $\geq 4 \times \text{HL-LHC}$
- Machine design within constraints from LHC CE and based on HL-LHC and FCC technologies  $10\text{ ab}^{-1}$  over 20 years.



# FCC-ee collider parameters

parameter	Z	WW	H (ZH)	ttbar
beam energy [GeV]	45	80	120	182.5
beam current [mA]	1390	147	29	5.4
no. bunches/beam	16640	2000	393	48
bunch intensity [ $10^{11}$ ]	1.7	1.5	1.5	2.3
SR energy loss / turn [GeV]	0.036	0.34	1.72	9.21
total RF voltage [GV]	0.1	0.44	2.0	10.9
long. damping time [turns]	1281	235	70	20
horizontal beta* [m]	0.15	0.2	0.3	1
vertical beta* [mm]	0.8	1	1	1.6
horiz. geometric emittance [nm]	0.27	0.28	0.63	1.46
vert. geom. emittance [pm]	1.0	1.7	1.3	2.9
bunch length with SR / BS [mm]	3.5 / 12.1	3.0 / 6.0	3.3 / 5.3	2.0 / 2.5
luminosity per IP [ $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ ]	>200	>25	>7	>1.4
beam lifetime rad Bhabha / BS [min]	68 / >200	49 / >1000	38 / 18	40 / 18





# FCC-ee operation model

working point	luminosity/IP [ $10^{34} \text{ cm}^{-2}\text{s}^{-1}$ ]	total luminosity (2 IPs)/ yr	physics goal	run time [years]
Z first 2 years	100	26 $\text{ab}^{-1}/\text{year}$	<b>150 <math>\text{ab}^{-1}</math></b>	<b>4</b>
Z later	200	52 $\text{ab}^{-1}/\text{year}$		
<i>W</i>	25	7 $\text{ab}^{-1}/\text{year}$	<b>10 <math>\text{ab}^{-1}</math></b>	<b>1-2</b>
<i>H</i>	7.0	1.8 $\text{ab}^{-1}/\text{year}$	<b>5 <math>\text{ab}^{-1}</math></b>	<b>3</b>
machine modification for RF installation & rearrangement: <b>1 year</b>				
top 1st year (350 GeV)	0.8	0.2 $\text{ab}^{-1}/\text{year}$	<b>0.2 <math>\text{ab}^{-1}</math></b>	<b>1</b>
top later (365 GeV)	1.4	0.36 $\text{ab}^{-1}/\text{year}$	<b>1.5 <math>\text{ab}^{-1}</math></b>	<b>4</b>

**total program duration: 14-15 years** - *including machine modifications*  
**phase 1 (Z, W, H): 8-9 years,**    **phase 2 (top): 6 years**





# FCC-pp collider parameters



parameter	FCC-hh		HE-LHC	HL-LHC	LHC
collision energy cms [TeV]	100		27	14	14
dipole field [T]	16		16	8.33	8.33
circumference [km]	97.75		26.7	26.7	26.7
beam current [A]	0.5		1.1	1.1	0.58
bunch intensity [ $10^{11}$ ]	1	1	2.2	2.2	1.15
bunch spacing [ns]	25	25	25	25	25
synchr. rad. power / ring [kW]	2400		101	7.3	3.6
SR power / length [W/m/ap.]	28.4		4.6	0.33	0.17
long. emit. damping time [h]	0.54		1.8	12.9	12.9
beta* [m]	1.1	0.3	0.45	0.15 (min.)	0.55
normalized emittance [ $\mu\text{m}$ ]	2.2		2.5	2.5	3.75
peak luminosity [ $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ ]	5	30	16	5 (lev.)	1
events/bunch crossing	170	1000	460	132	27
stored energy/beam [GJ]	8.4		1.3	0.7	0.36

# FCC-pp layout






# Worldwide FCC Nb<sub>3</sub>Sn program



Main development goal is wire performance increase:

- $J_c$  (16T, 4.2K) > 1500 A/mm<sup>2</sup> → 50% increase wrt HL-LHC wire
- Reduction of coil & magnet cross-section

After only one year of development, **prototype Nb<sub>3</sub>Sn wires from several new industrial FCC partners already achieve HL-LHC performance**




Conductor activities for FCC started in 2017:

- Bochvar Institute (production at TVEL), Russia
- KEK (Jastec and Furukawa), Japan
- KAT, Korea
- Columbus, Italy
- University of Geneva, Switzerland
- Technical University of Vienna, Austria
- SPIN, Italy
- University of Freiberg, Germany

In addition, agreements under preparation:

- Bruker, Germany
- Luvata Pori, Finland






# 16 T dipole design activities and options




Cos-theta



INFN




Blocks



CEA

Common coils




CIEMAT

Swiss contribution



Canted Cos-theta





PSI



The U.S. Magnet Development Program Plan




S. A. Gourlay, S. O. Prestemon  
Lawrence Berkeley National Laboratory  
Berkeley, CA 94720  
A. V. Zlobin, L. Cooley  
Fermi National Accelerator Laboratory  
Batavia, IL 60510  
D. Larbalastier  
Florida State University and the  
National High Magnetic Field Laboratory  
Tallahassee, FL 32310



LBNL

FNAL




Short model magnets (1.5 m lengths) will be built from 2018 – 2022  
Russian 16 T magnet program launched by BINP recently.





# Technical Schedule for each of the 3 options



**schedule constrained by 16 T magnets & CE**  
→ earliest possible beam operation dates

- FCC-ee: 2039
- FCC-hh: 2043
- HE-LHC: 2040 (with HL-LHC stop LS5 / 2034)



# Global FCC Collaboration



124

Institutes

30


Companies

32

Countries




# LHeC



W Kandinsky, Circles in a Circle, 1923, Philadelphia Art Museum

Slides from Max Klein for LHeC UK



# Three Raisons d'être of the LHeC

## Physics

- **Microscope:** World's Cleanest High Resolution
- **Maximises** the LHC Physics Programme
- **Creation** of a high precision, novel Higgs facility
- **Discovery** Beyond the Standard Model
- **Revolution** of Nuclear Particle Physics

## Technology

**Accelerator:** Novel SRF ERL, green power facility  
**Detector:** Novel high tech (CMOS..) apparatus

→ Keep accelerator and detector base uptodate while preparing for colliders that cost  $O(10)$ BSF

## Sustainability and Cost

### LHC:

- see: SM, Higgs and no BSM
- use: Investment of  $O(5)$  BSF
- run: HL LHC until ~2040

**LHeC** [1206.2913, update 2/19]

- 1.2 TeV ep/A for  $O(1)$ BSF


→ Establish novel ep+pp

**Twin Collider Facility at CERN:**



sustains HL LHC and bridges to  
 CERN's long term future

**For installation during LS4 (2030+)  
 and long term use (HE LHC, FCCeh)**

# Prospects for Higgs Couplings: LHC, LHeC, ep+pp, ILC



HL-LHC ATLAS2014. LHeC:  $1ab^{-1}$ , in progress ep+A14 ILC:  $2ab^{-1}$ : 1710.0761  
 HL LHC WS to come, ep: J de Blas, M+U Klein ep+pp: J de Blas, prel.  $e^+e^-$  provides  $\Gamma_{tot}$




## Physics

1802.04317

$\alpha_s$  to 0.1%  
 $V_{tb}$  to 1%...  
 $\sin^2\theta_w$  better than LEP  
 $M_w$  [pp+ep]: 0.007%  
 ...  
 HIGH precision leads  
 to Discovery BSM


PPAP 16/7/18 -- I. Shipsey






- 2 Linacs (Four 5-Cell 801.58 MHz SC cavities)
- 3 turns (160 MeV/turn)
- Max. beam energy 500 MeV

## PERLE at Orsay




Challenge: demonstrate multi-turn ERL (cbeta, 2019)  
Develop 802 MHz, LHeC Technology (PERLE > 2022)




**PERLE/LHeC first 802 MHz cavity**  
CERN-Jlab.  $Q_0 > 3 \cdot 10^{10}$ , 2K. freedom  
to choose operation point  $\sim 20$  MV/m

PPAP 16/7/18 -- I. Shipsey

# Acc & Det Technology



Zoom LHeC detector [ $15.6 \times 10.4 \text{ m}^2$  HE LHC]



## UK Institutes

### Accelerator

AsTEC, Cockcroft (Lancaster, Manchester, Liverpool, Strathclyde), JAI (Oxford)

### Detector+Physics

Birmingham, Liverpool, Manchester, Oxford, QMW

HERA+LHC have also  
Bristol, Glasgow, Imperial, Lancaster, RAL, UCL.

Detector: a new task post HL LHC design  
Challenge for Acc+Det: 3 beam-IR design

Most up-to-date Information: <https://indico.cern.ch/event/698368/>

Workshop: LHeC/FCCeh and PERLE  
Two week ago at Orsay near Paris



**Electrons for the LHC**  
LHeC/FCCeh and PERLE  
Workshop

**June 27-29, 2018**  
LAL-Orsay, France

**Organising Committee:**  
Nestor Armesto (USC)  
Oliver Brüning (CERN)  
Walid Kaabi (LAL)  
Uta Klein (Liverpool)  
Zhiqing Zhang (LAL)

**Advisory Committee:**  
Sergio Bertolucci (Bologna)  
Nicola Bianchi (INFN)  
Frederick Bordry (CERN)  
Oliver Brüning (CERN)  
Stanley Brodsky (SLAC)  
Weiheng Chen (IHEP Beijing)  
Eckhard Eiten (CERN)  
Stefano Forte (Milano)  
Andrew Hutton (Jefferson Lab)

**Coordination Group:**  
Nestor Armesto (Garriga de Compostela)  
Gianluigi Arduini (CERN)  
Oliver Brüning (CERN)  
Andreas Gaidi (CERN)  
Erik Jensen (CERN)  
Walid Kaabi (LAL Orsay)  
Max Klein (Liverpool)  
Peter Koslowski (Liverpool)

**Physics Convenors:**  
Hassan Arora (Garriga de Compostela)  
Georgios Antoniadis (CERN)  
Maurizio Basso (CERN)  
David Fuster (CERN)  
David Goussard (CERN)  
Dimitris Karas (CERN)  
Paul Marage (CERN)  
Paul Marage (Birmingham)  
Daniel Schulte (CERN)  
Frank Zimmermann (CERN)

**Logos:** PERLE, FCC, LHC, CERN, INFN, SLAC, IHEP, ARIES, APEC, DESY

<https://indico.cern.ch/event/698368/>



### New and Updates on

**Physics:** PDFs, QCD, H, t, BSM, eA + Relation eh-hh..

**Accelerator:** IR, Optics, Lattice, Cost-Energy, CE..

**Detector:** the GPD and its fwd and bwd detectors


**PERLE:** Source, Injector, Cavity, Cryomodule,.. Physics  
**Project Development** towards the ES2020:

LHeC + FCCeh+ PERLE input 12/18. **PERLE TDR in 2019.**

<http://lhec.web.cern.ch>



# Circular electron positron Collider (CEPC)



April 2017

June 2018


Lumi.	Higgs	W	Z	Z(2T)
$\times 10^{34}$	2.93	11.5	16.6	32.1

- ✓ double ring baseline design
- ✓ switchable between H and Z/W w/o hardware change (magnet switch)
- ✓ use half SRF for Z and W
- ✓ can be optimized for Z with 2T detector (~3200× LEP luminosity)


Intl. review - June 28-30 at IHEP

Release of CDR: July (accelerator), September (detector)

# CEPC Accelerator Chain and Systems



**Booster Cycle (0.1 Hz)**




**The key systems of CEPC:**

- 1) Linac Injector
- 2) Booster
- 3) Collider ring
- 4) Machine Detector Interface
- 5) Civil Engineering

**CDR provides details of all systems**

# The 100k tunnel cross section




**SPPC**  
collider

**CEPC**  
collider

**CEPC**  
Booster

**CEPC Civil Engineering Design very advanced**

**Proposed in Lausanne Workshop in 1984**




**LEP tunnel internal diameter is 3.8 metres in the arcs  
4.4 or 5.5 metres in the straight sections**




# Tunnel cross sections for HE-LHC, SppC and FCC-hh


**FCC-hh**




**SppC**



**HE-LHC**



# The CEPC Baseline Collider Design




**Double ring**

**Common RF cavities for Higgs**

**Two RF sections in total**

**Two RF stations per RF section**

**$10 \times 2 = 20$  cryomodules**



**6 2-cell cavities per cryomodule**



# Site selection

**Chuangchun, Jilin**  
吉林长春




**Huangling, Shanxi**  
陕西黄陵



## Considerations:

1. Available land
2. Geological conditions
3. Good social, environment, transportation and cultural conditions
4. Fit local development plan: mid-size city → + science city

**Shenshan, Guangdong**  
深汕合作区



**Qinhuangdao, Hebei**  
河北秦皇岛



**Xiong an, Hebei**  
河北雄安




**Huzhou, Zhejiang**  
浙江湖州







# Detector Conceptual Designs (CDR)

Baseline detector (3 Tesla)  
ILD-like  
(similar to pre-CDR)



Low  
magnetic field  
concept  
(2 Tesla)



Full silicon  
tracker  
concept


Final **two** detectors likely to be a mix and match of different options

# CEPC Accelerator CDR Completed


CEPC accelerator CDR **completed** in June 2018 (to be printed in **July 2018**)

## ➔ Executive Summary


1. Introduction
  2. Machine Layout and Performance
  3. Operation Scenarios
  4. CEPC Booster
  5. CEPC Linac
  6. Systems Common to the CEPC Linac, Booster and Collider
  7. Super Proton Proton Collider
  8. Conventional Facilities
  9. Environment, Health and Safety
  10. R&D Program
  11. Project Plan, Cost and Schedule
- Appendix 1: CEPC Parameter List
- Appendix 2: CEPC Technical Component List
- Appendix 3: CEPC Electric Power Requirement
- Appendix 4: Operation for High Intensity  $\gamma$ -ray Source
- Appendix 5: Advanced Partial Double Ring
- Appendix 6: CEPC Injector Based on Plasma Wakefield Accelerator
- Appendix 7: International Review Report



**March 2015**



**April 2017**




**July 2018**

**Physics and Detector CDR  
to follow soon afterwards  
(Need to adapt to recent modifications)**



# CEPC “optimistic” Schedule



# Current CEPC Organization

Since Sept.  
2013



## Institutional Board

YN GAO  
J. GAO

## Steering Committee

Y.F. WANG (IHEP),....

## Project Director

XC LOU  
Q. QIN  
N. XU

## Theory

HJ HE(TH)  
JP MA(ITP)  
XG HE(SJTU)

## Accelerator

J. GAO (IHEP)  
CY Long (IHEP)  
SN FU (IHEP)

## Detector

Joao Costa (IHEP)  
S. JIN (NJU)  
YN GAO (TH)

## International Advisory Committee

Young-Kee Kim, U. Chicago (Chair)  
Barry Barish, Caltech  
Hesheng Chen, IHEP  
Michael Davier, LAL  
Brian Foster, Oxford  
Rohini Godbole, CHEP, Indian Institute of Science  
David Gross, UC Santa Barbara  
George Hou, Taiwan U.  
Peter Jenni, CERN  
Eugene Levichev, BINP  
Lucie Linssen, CERN  
Joe Lykken, Fermilab  
Luciano Maiani, Sapienza University of Rome  
Michelangelo Mangano, CERN  
Hitoshi Murayama, UC Berkeley/IPMU  
Katsunobu Oide, KEK  
Robert Palmer, BNL  
John Seeman, SLAC  
Ian Shipsey, Oxford  
Steinar Stapnes, CERN  
Geoffrey Taylor, U. Melbourne  
Henry Tye, IAS, HKUST  
Yifang Wang, IHEP  
Harry Weerts, ANL



# CEPC meetings and international impact

Many international events have been hosted to discuss CEPC physics and carry out collaboration on key-technology research

Next workshop  
April 2019 in Oxford

## INTERNATIONAL WORKSHOP ON HIGH ENERGY CIRCULAR ELECTRON POSITRON COLLIDER

November 6-8, 2017  
IHEP, Beijing

<http://indico.ihep.ac.cn/event/6618>

### International Advisory Committee

Young-Kee Kim, U. Chicago (Chair)  
Barry Barish, Caltech  
Hesheng Chen, IHEP  
Michael Davier, LAL  
Brian Foster, Oxford  
Rohini Godbole, CHEP, Indian Institute of Science  
David Gross, UC Santa Barbara  
George Hou, Taiwan U.  
Peter Jenni, CERN  
Eugene Levichev, BINP  
Lucie Linssen, CERN  
Joe Lykken, Fermilab  
Luciano Maiani, Sapienza University of Rome  
Michelangelo Mangano, CERN  
Hitoshi Murayama, UC Berkeley/PMU  
Katsunobu Oide, KEK  
Robert Palmer, BNL  
John Seeman, SLAC  
Ian Shipsey, Oxford  
Steinar Stapnes, CERN  
Geoffrey Taylor, U. Melbourne  
Hiroaki Teraoka, IHEP  
Yuanji Wang, IHEP  
Yuanji Wang, IHEP

### Local Organizing Committee

Xinchou Lou, IHEP (Chair)  
Qinghong Cao, PKU  
Joao Guimaraes Costa, IHEP  
Jie Gao, IHEP  
Yuanming Gao, THU  
Hongjian He, THU  
Shan Jin, IHEP  
Gang Li, IHEP  
Jianbei Liu, USTC  
Yajun Mao, PKU  
Qing Qin, IHEP  
Manqi Ruan, IHEP  
Meng Wang, SDU  
Nu Xu, CCNU  
Haijun Yang, SJTU  
Hongbo Zhu, IHEP

260 attendees  
30% from foreign institutions



## Workshop on the Circular Electron-Positron Collider

EU Edition

Roma, May 24-26 2018  
University of Roma Tre



55% attendance from abroad

<https://agenda.infn.it/conferenceDisplay.py?ovw=True&confid=14816>

### Scientific Committee

Franco Bedeschi - INFN, Italy  
Alain Blondel - Geneva Univ., Switzerland  
Daniela Bortoletto - Oxford Univ., UK  
Manuela Boscolo - INFN, Italy  
Biagio Di Micco - Roma Tre Univ. & INFN, Italy  
Yunlong Chi - IHEP, China  
Marcel Demarteau - ANL, USA  
Yuanming Gao - Tsinghua Univ., China  
Joao Guimaraes da Costa - IHEP, China  
Gao Jie - IHEP, China  
Gang Li - IHEP, China  
Jianbei Liu - USTC, China  
Xinchou Lou - IHEP, China  
Felix Sefkow - DESY, Germany  
Shan Jin - Nanjing Univ., China  
Marcel Vos - CSIC, Spain

### Local Organizing Committee

Antonio Baroncelli - INFN, Italy  
Biagio Di Micco - Roma Tre Univ. & INFN, Italy  
Ada Farilla - INFN, Italy  
Francesca Paolucci - Roma Tre Univ. & INFN, Italy  
Domizia Orestano - Roma Tre Univ. & INFN, Italy  
Marco Sessa - Roma Tre Univ. & INFN, Italy  
Monica Verducci - Roma Tre Univ. & INFN, Italy




- \* The accelerator CDR has been completed satisfying the luminosity requirements both as a Higgs and Z factory
  - \* Detector CDR will follow soon
- \* Key technologies are under R&D and put to prototyping:
  - \* Accelerator: SC cavity, high efficiency klystron, low field precision magnet, copper vacuum chamber, HTS, ...
  - \* Detector: Pixel detector, TPC, PFA-based electromagnetic and hadronic calorimeters, magnet, ...
- \* CEPC civil engineering design and site selection going well
- \* CEPC funding adequate for required R&D program
- \* CEPC interest abroad is steadily increasing
- \* From 2018–2022, CEPC TDR will be finished with accelerator key hardware R&D completed and industrialization ready for construction start in 2022



# Future Plan of IHEP

Part 3





# IHEP Large Science Facilities

Part 1



## In operation (6)

1. Beijing Electron Positron Collider (BEPCII) / (BESIII)
2. Beijing Synchrotron Radiation Facility (BSRF)
3. Yangbajing Cosmic Ray Observatory (YBJ)
4. Daya Bay Reactor Neutrino Experiment
5. Hard X-ray Modulation Telescope (HXMT)
6. China Spallation Neutron Source (CSNS)

## Under construction (4)

1. Jiangmen Underground Neutrino Observatory (JUNO)
2. Large High-Altitude Air Shower Observatory (LHAASO)
3. Ali CMB Polarization Telescope (AliCPT-1)
4. High Energy Photon Source (HEPS/HEPS-TF)


## Under planning (6)

1. China Initiative Accelerator Driven System (CiADS)
2. China Spallation Neutron Source II (CSNSII)
3. Enhanced X-ray Timing and Polarimetry mission (eXTP)
4. High Energy cosmic-Radiation Detection (HERD)
5. Circular Electron Positron Collider-Super proton-proton collider (CEPC-SppC)
6. Other Light Source Projects : Southern Photon Source 、 SCLS.....

# Accelerator R&D – Advanced Novel Accelerators (ICFA Panel)



**ALEGRO** (Advanced LinEar collider study GROup, for a multi-TeV Advanced Linear Collider) Workshop (March 2018 in Oxford): <http://www.physics.ox.ac.uk/confs/alegro2018/index.asp>

The objective of this first ALEGRO workshop was to prepare and deliver, by the end of 2018, a document detailing the international roadmap and strategy of Advanced Novel Accelerators (ANAs) with clear priorities as input for the European Particle Physics Strategy Update.



# Current UK advanced accelerator technique work

Technique/beam	Groups/facilities
Laser driven (LWFA) electrons	CI (Lan, Liv, Man, Str) – CLF/SCAPA JAI (Imp, Oxf) - CLF
Laser driven (LWFA) positrons	QUB - CLF
Laser driven (LWFA) protons/ions	Imp, Str, QUB, York – CLF/SCAPA
Electron driven (PWFA) electrons	CI (Lan, Liv, Man, Str) – CLARA/FACET
Proton driven (PWFA) electrons	Lan, Liv, Man, UCL – AWAKE



PWFA



UK Plasma Wakefield Accelerator Roadmap 2018

## UK roadmap for plasma acceleration research

"Although, having an electron-positron linear collider as an ultimate aim for plasma wakefield acceleration and working towards achieving a number of its parameters is very valuable, it is prudent to consider other first applications.

One should however distinguish between the first plasma acceleration application to a stand-alone all plasma acceleration collider, and an upgrade of conventional collider with plasma acceleration. While it is at this moment inconceivable to suggest an all-plasma electron-positron collider, it is reasonable to consider plasma acceleration upgrade for either ILC or CLIC colliders, in case if construction of the first Higgs-Factory or Top-Factory stage of either of them will be approved. Given the rate of the progress of plasma acceleration technology, it is entirely possible to consider their upgrades to TeV energy using plasma acceleration."

<http://pwasc.org.uk/uk-roadmap-development>