

Quantum Information and Entanglement with Top Quarks at the LHC

TOP 2020

Based on: [2003.02280](#)


Y. Afik, J. R. M. De Nova

16.09.2020



Motivation

- The Standard Model is a Quantum Field Theory: Special Relativity and Quantum Mechanics.
- Fundamental properties of Quantum Mechanics can be tested via the Standard Model.
- Entanglement is one of the most genuine features of Quantum Mechanics.
- First study of entanglement between a pair of quarks.
- Applying quantum tomography, which is a genuine Quantum Information technique.
- Quantum Information \rightarrow High Energy physics.




Top-Quark

- **General:**

- Hadronisation: $\sim 10^{-23}\text{s}$.
- Spin-decorrelation: $\sim 10^{-21}\text{s}$.

- **Top quark:**

- Lifetime: $\sim 10^{-25}\text{s}$.
- Spin information \rightarrow decay products.
- Spin-correlations between a pair of top-quarks can be measured.
- Considering leptonic decays.




Spin-Correlations between Top-Quark Pairs

- Studied extensively theoretically.
- Measured by the D0, CDF, ATLAS and CMS collaborations.
- No link between spin-correlations and quantum entanglement so far.
- Note! **Spin-Correlations** \neq **Quantum Entanglement!** However, Quantum Entanglement \subset Spin-Correlations.



LO Analytical Calculation



- Analytical calculation at LO.


- The total quantum state:

$$\rho(M_{t\bar{t}}) \equiv \int_{2m_t}^{M_{t\bar{t}}} dM \int d\Omega \rho(M, \hat{k}) \rho(M, \hat{k})$$



Entanglement Criterion

- **Concurrence** $C[\rho]$: Quantitative measurement of entanglement.
- $0 \leq C[\rho] \leq 1$, $C[\rho] \neq 0$ iff the state is entangled.
- Here, $C[\rho] = \max(\Delta, 0)/2$. Entanglement equivalent to $\Delta > 0$.
- $D = \frac{\text{tr}[\mathbf{C}]}{3} = -\frac{1+\Delta}{3}$ provides an experimental entanglement marker.



Measurable Entanglement Marker

- Plots are shown with integration only for $[2m_t, M_{t\bar{t}}]$.
- In particular:
$$\frac{1}{\sigma} \frac{d\sigma}{d\cos\varphi} = \frac{1}{2}(1 - D \cos\varphi)$$
 where φ is the angle between the lepton directions in each one of the parent top and antitop rest frames.
- The condition $\Delta > 0$ translates into $D < -1/3$.




Figure: Up: concurrence; bottom: Statistical deviation from the null hypothesis ($D = -1/3$).

Quantum Tomography

- Quantum tomography=Reconstruction of the quantum state of the system.
- Only need to measure 4 parameters (transverse and longitudinal spin correlations and the longitudinal polarizations).
- Test theoretical predictions for the $t\bar{t}$ quantum state.



Summary

- First study of entanglement between quarks.
- Quantum Information study in a relativistic system.
- Although the calculation is analytical at leading order, the conclusion still holds at higher orders.
- Quantum tomography: new platform to test new theories and new physics effects.
- Interdisciplinary measurement: propagate Quantum Information theory into High Energy physics.

Summary

- First study of entanglement between quarks.
- Quantum Information study in a relativistic system.
- Although the calculation is analytical at leading order, the conclusion still holds at higher orders.
- Quantum tomography: new platform to test new theories and new physics effects.
- Interdisciplinary measurement: propagate Quantum Information theory into High Energy physics.
- **Can be detected at the LHC with currently recorded data!**

Thank You



THANK
YOU!

The background of the slide features a soft, ethereal glow with two prominent, stylized atomic models. Each model consists of a central nucleus of yellow and orange particles, surrounded by a translucent blue sphere representing the electron cloud. Inside these spheres, thin yellow lines and small white dots suggest the paths and positions of electrons. A bright, multi-colored beam of light, transitioning from purple to yellow, originates from the left and points towards the right, passing behind the word 'Backup'.

Backup

NLO Corrections

- LO: analytical calculation.
- NLO: numerical calculation by using Monte Carlo simulation.
- MADGRAPH and MADSPIN are used.
- Good agreement is observed.





Figure: The value of D within the mass window $[2m_t, M_{t\bar{t}}]$. The horizontal line represents the critical value $D = -1/3$.

Local Realism

- Locality: physical influences do not propagate faster than light.
- Realism: physical properties are defined before, and independent of observation.
- Both of the assumptions (together, not separately) are in tension with Quantum Mechanics.



Top-Quark Pair Spin Density Matrix


- General form:

$$\rho = \frac{I_4 + \sum_i (B_i^+ \sigma^i + B_i^- \bar{\sigma}^i) + \sum_{i,j} C_{ij} \sigma^i \bar{\sigma}^j}{4}$$

- $\sigma^i/2, \bar{\sigma}^i/2$ - spin operators of the top, antitop.
- B_i^+, B_i^- characterize the spin polarizations, $B_i^+ = \langle \sigma^i \rangle$, $B_i^- = \langle \bar{\sigma}^i \rangle$.
- At LO $B_i^\pm = 0$.
- C_{ij} the $t\bar{t}$ spin correlations, $C_{ij} = \langle \sigma^i \bar{\sigma}^j \rangle$.

What is Quantum Entanglement?


- Quantum state of one particle cannot be described independently from another particle.
- \Rightarrow **Correlations** of observed physical properties of both systems.
- \Rightarrow **Measurement** performed on one system seems to be instantaneously influencing other systems entangled with it.




- Observed in photons, atoms, superconductors, mesons, analog Hawking radiation, nitrogen-vacancy centers in diamond and even macroscopic diamond.

High Energy Physics Example

- At B-Factories, e^+e^- collisions can be properly adjusted in order to create $\Upsilon(4S)(b\bar{b})$.
- $\Upsilon(4S)(b\bar{b})$ decays to $B^0 + \bar{B}^0$, where we have $|B^0\rangle = |\bar{b}d\rangle, |\bar{B}^0\rangle = |b\bar{d}\rangle$.
- We get an entangled state:
$$\frac{1}{\sqrt{2}}(|B^0\rangle|\bar{B}^0\rangle - |\bar{B}^0\rangle|B^0\rangle).$$



EPR Paradox



A. Einstein

B. Podolsky

N. Rosen

MAY 15, 1935

PHYSICAL REVIEW

VOLUME 47


Can Quantum-Mechanical Description of Physical Reality Be Considered Complete?

A. EINSTEIN, B. PODOLSKY AND N. ROSEN, *Institute for Advanced Study, Princeton, New Jersey*

(Received March 25, 1935)

EPR Paradox

- Entanglement: "spooky action at a distance" (A. Einstein).




- Assuming two particles with spacial distance.
- When a measurement is done on one of the particles, the other one "knows" about it immediately.
- Information travel faster than light?
- Contradicts the theory of relativity.
- **Conclusion:** the theory of Quantum Mechanics is incomplete.

Hidden Variables

- By EPR, each particle "carries" variables that knows the state before the measurement.
- \Rightarrow There are some hidden variables that are missing in order to have a full theory.
- The Copenhagen Interpretation: superposition of states until a measurement was done.
- Bohr Vs. Einstein.

"God does not play at dice with the universe".



"Quit telling God what to do!"

- Who is right?



III.5 ON THE EINSTEIN PODOLSKY ROSEN PARADOX*

JOHN S. BELL†

- If local hidden variables holds, they should satisfy some inequality.
- $C(x, y)$ are the correlations between different measurements at different detectors.
- The parameters a, b, c are different directions for the measurement.
- Original form: $1 + C(b, c) \geq |C(a, b) - C(a, c)|$.

Quantum State

- **Pure state:** can be described by wave-functions $\sum_i \alpha_i \cdot |\psi_i\rangle$.
- **Mixed state:** can be described by a density matrix: $\rho = \sum_i p_i \cdot |\psi_i\rangle \langle \psi_i|$.
 - Example: at the LHC we cannot control the initial state.
- Some inequalities can be measured related to ρ , providing an entanglement criterion.



Mathematical Formalism

- Two different systems A and B: $\mathcal{H} = \mathcal{H}_a \otimes \mathcal{H}_b$.
- Separable: $\rho = \sum_n p_n \rho_n^a \otimes \rho_n^b$.
- $\rho_n^{a,b}$ are quantum states in A, B, $\sum_n p_n = 1$, $p_n \geq 0$
- Classically correlated state in $\mathcal{H} \rightarrow$ can be written in this form.
- Non-separable state is called entangled and hence, it is a non-classical state.




Separable



Non-Separable

Intuition: Spin States at Threshold

- The state is determined by the initial spins.
- $q\bar{q}$: $\rho^{q\bar{q}} = (|\uparrow_{\hat{p}}\uparrow_{\hat{p}}\rangle \langle\uparrow_{\hat{p}}\uparrow_{\hat{p}}| + |\downarrow_{\hat{p}}\downarrow_{\hat{p}}\rangle \langle\downarrow_{\hat{p}}\downarrow_{\hat{p}}|) / 2$.
- gg : $\rho^{gg} = |\Psi_0\rangle \langle\Psi_0|$, with $|\Psi_0\rangle = (|\uparrow_{\hat{p}}\downarrow_{\hat{p}}\rangle - |\downarrow_{\hat{p}}\uparrow_{\hat{p}}\rangle) / \sqrt{2}$.
- $q\bar{q} \rightarrow$ correlated, not entangled; $gg \rightarrow$ correlated, entangled.



Basis Selection

- Helicity basis: $\{\hat{k}, \hat{r}, \hat{n}\}$:
 - \hat{k} - direction of the top in the $t\bar{t}$ CM frame.
 - \hat{p} - direction of the beam.
 - $\cos \Theta = \hat{k} \cdot \hat{p}$.
 - $\hat{r} = (\hat{p} - \cos \Theta \hat{k}) / \sin \Theta$.
 - $\hat{n} = \hat{r} \times \hat{k}$.
 - Describe each individual process with a **fixed direction**.
- Beam basis: $\{\hat{x}, \hat{y}, \hat{z}\}$:
 - \hat{z} along the beam axis.
 - \hat{x}, \hat{y} transverse directions to the beam.
 - After averaging: $C_x = C_y = C_{\perp}$.
 - Studying the **total quantum state**.






Figure: Helicity basis.
Figure is from [Phys. Rev. D 100, 072002](#).

Spin-Correlations - $M_{t\bar{t}}$ Dependence

- Plots are shown with integration only for $[2m_t, M_{t\bar{t}}]$.
- c) Spin-correlations after $[2m_t, M_{t\bar{t}}]$ integration.
Main: C_\perp , C_z , D (quantum tomography).
Inset: C_{rr} , C_{nn} , C_{kk} .
- d) Main plot: Concurrence.
Inset: integrated cross-section.




Entanglement Criterion - Concurrence

- Concurrence:

$$C[\rho] \equiv \max(0, \lambda_1 - \lambda_2 - \lambda_3 - \lambda_4)$$

- λ_i are the eigenvalues of the Concurrence matrix $\mathcal{C}(\rho)$.
- In our case $\mathcal{C}(\rho) = \rho$.
- $0 \leq C[\rho] \leq 1$, vanishing if and only if the state is separable.
- Compute the eigenvalues of ρ - apply a criterion for entanglement.



Entanglement Criterion - Peres-Horodecki

- Partial transpose in one subsystem. Example:

$$\rho^{T_B} = \sum_n p_n \rho_n^a \otimes (\rho_n^b)^T$$

- If ρ is separable, all of the eigenvalues of ρ^{T_B} are non-negative.
- Reduces to the condition $\Delta > 0$, with $\Delta \equiv -C_{nn} + |C_{kk} + C_{rr}| - 1$.
- Total quantum state: can use any orthonormal basis to characterize entanglement.
- Link to concurrence: $C = \max(\Delta, 0)/2$.
- We also depict: $D = \frac{\text{tr}[C]}{3} = -\frac{1+\Delta}{3}$.

